



Effect of Stocking Density on Growth Performance and the Survival of Golden Mahseer, *Tor putitora* (Hamilton) Fry

Md. Golam Sajed Riar^{1*}, Nur- A Raushon² and Sumit Kumer Paul³

¹Head Quarter, Bangladesh Fisheries Research Institute, Bangladesh.

²Freshwater Station, Bangladesh Fisheries Research Institute, Bangladesh.

³Department of Fisheries, Bangamata Sheikh Fojilatunnesa Mujib Science and Technology University, Jamalpur, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2021/v14i530308

Editor(s):

- (1) Prof. Ahmed Karmaoui, University Moulay Ismail, Morocco.
- (2) Dr. Luis Enrique Ibarra Morales, University of Sonora, Mexico.
- (3) Dr. Pinar Oguzhan Yildiz, Ataturk University, Turkey.
- (4) Dr. Telat Yanik, Atatürk University, Turkey.

Reviewers:

- (1) Mohamed Fathy Aid Abdel-Aziz, Egypt.
- (2) Mohamed Ahmed Aly El-Drawany, Zagazig University, Egypt.

Complete Peer review History: <https://www.sdiarticle4.com/review-history/73736>

Original Research Article

Received 04 July 2021
Accepted 14 September 2021
Published 22 September 2021

ABSTRACT

Growth performance and survival of *Tor putitora* fry under different stocking densities were evaluated fry rearing system. The experimental period was 10 weeks from 31 December to 10 March 2019. The experiment was carried out in nine earthen ponds of 0.04 ha each under three treatments with three replications. Thirty days old fry were stocked at the rate of $1.0 \times 10^5/\text{ha}$ was designated as treatment- 1 (T_1), $1.5 \times 10^5/\text{ha}$ as treatment- 2 (T_2) and $2.0 \times 10^5/\text{ha}$ as treatment- 3 (T_3), respectively. Fry were fed with commercially available nursery feed containing 32% crude protein. It was observed that, lower stocking density showed highest daily weight gain in T_1 (growth 0.066 ± 0.006 g/day) compared with higher stocking density in T_2 (growth 0.044 ± 0.004 g/day) and T_3 (growth 0.024 ± 0.003 g/day), respectively. It is also noticed that, the lower stocking showed the highest survival rate (79.66 ± 4.34 %) than the other two treatments (66.97 ± 3.67 and 54.67 ± 3.12 %). The values of different water quality parameters were within the optimum ranges for the rearing

*Corresponding author: Email: riarsajed@yahoo.com, riarsajed@gmail.com;

of carp fry. Water quality parameters did not show significant variations in the experimental ponds under different stocking densities. Among these three stocking densities lower stocking density (T_1) showed the best result compare with the other two higher stocking densities.

Keywords: *Tor putitora*; nursing system; stocking density; growth performance; survival.

1. INTRODUCTION

Golden Mahseer (*Tor putitora*) is one of the most important endangered fish species of Bangladesh. Once the fish was widely distributed throughout Bangladesh, India, Nepal and Pakistan, considered as a popular sports fish for this region [1]. This fish is highly popular for its qualities and test. This fish contribute a significant amount in total fish production of The Indian subcontinent [2].

Tor genus consists of more than 20 other species which are not yet scientifically identified [3]. Two species of these, *Tor tor* and *Tor putitora* are available in the hilly streams of Sylhet, Mymensingh, Dinajpur and Chittagong districts in Bangladesh [4]. The fish is habituated with a wide range of temperature, from semi-cold waters of foot hills to warm waters. The body is stout and bearing medium to large size barbs and attains nearly 3 m in length and 45 kg in weight [5]. By nature it is omnivorous but in early stage of life, it show carnivore feeding habit and feed on zooplankton, other fish larvae, insects etc. adult *Tor* mainly feed on other fish larvae, small mollusks and algae grow on rocks [6]. *Tor* species prefer sandy ground with small rocks for breeding with high oxygen and comparative higher oxygen from other parts of its habitat [7]. November to January is the main breeding season of this species in Bangladesh [8]. *T. putitora* is a low fecund fish compared with other carp species and its fecundity ranges between 8000 to 12,000 eggs/kg [9].

Due to prolonged drought, frequent devastating floods, siltation and soil erosion in the hilly rivers, reservoirs and man-made changes i.e. construction of flood control measures and drainage structures, dumping of agrochemicals and industrial pollutants, indiscriminate and destructive fishing practices in the aquatic ecosystem the natural stocks of Mahseer fishes have declined. These calamities damage the natural breeding ground and also destroy the nursing ground, feeding habitat; in a word destroy total population of this species [9]. This fish are now enlisted as a critically endangered species [10]. Now it's urgent needed to protect

this fish from declining. Not only the habitat degradation natural habitat, also the non-availability of fry and spawn due to improper nursery management system also an important cause of declining. This fish has also great potential and high demand for polyculture with other carps. But due to in sufficient supply of fry the culture system didn't develop properly. Bangladesh fisheries research Institute working on this fish from many years and develop its breeding protocol, single stage nursery system [9,7,8]. But to develop proper nursery maangemnet system it is also essential to develop fry culture system or secondary nursing system. Growth, survival and production of spawn and fry in ponds depend on stocking density, type and quantity of fertilizers and supplementary feeds. Therefore, the present study was undertaken to assess the effects of stocking density on *T. putitora* fry under controlled condition.

2. MATERIALS AND METHODS

2.1 Site Selection and Experimental Design

The experiment was conducted in the Muktagacha sub district, Mymensingh district, Bangladesh, during the period from 31 December to 10 March 2019 (70 days). Nine earthen ponds of 0.04 ha (40 X 10 m) with 1 m depth were selected for the experiment. Three treatments with three replications were considered under complete randomized design (CRD) for this experiment. Thirty days old fry were stocked at the rate of $1.0 \times 10^5/\text{ha}$ as treatment- 1 (T_1), $1.5 \times 10^5/\text{ha}$ as treatment- 2 (T_2) and $2.0 \times 10^5/\text{ha}$ as treatment- 3 (T_3), respectively with three replications each.

2.2 Pond Preparation, Stocking and Management

The ponds were prepared following standard management practices [11,4,6]. Ponds having proper inlet and outlet facilities. These were cleaned and dry under sunlight to remove aquatic vegetation and all unwanted species.

After drying 250 kg/ha lime was applied over the pond bottom. And after seven days of liming, ponds were filled with groundwater at 1m depth. Then 25 kg/ha urea and 50 kg/ha TSP were applied to produce the live feed for the spawn. Seven days after fertilization a fine mesh net used to remove harmful insect and predatory zooplankton. Thirty days old spawn of *T. putitora* obtained from Bangladesh Fisheries Research Institute (BFRI) hatchery were stocked in experimental ponds (Average length 3.12 ± 0.65 cm and average weight 1.12 ± 0.12 g) according to the experimental design. Spawn was fed with commercial feed two times daily at the early morning and evening. For 1st week 10% feed of total body weight was applied while the amount of feed increasing continuously but the percentage of feed decreasing 1% per week. Amount of feed adjust with weekly sampling of experimental fish. Moreover, ponds were fertilized every seven days with 12.5 kg/ha urea and 25 kg/ha TSP respectively.

2.3 Proximate Composition of Feed

Same feed were used in all treatments of this experiments. Commercially available nursery fish feed was purchased from the commercial Feed company, Bangladesh. Proximate compositions of the feeds (moisture, protein, lipid, ash and fiber) were analyzed [12]. The approximate composition of the feeding is given as shown in Table 1.

2.4 Water Quality Parameters

The important physio-chemical parameters of water viz., water temperature, pH, dissolved oxygen (DO), total alkalinity and ammonia were analyzed at every seven days interval to ensure that water quality was within permissible limits. The samples for the study were collected between 08:00 and 09:00 h.

2.5 Fish Sampling

Spawn was collected weekly with fine-meshed net for growth analysis. At the 71 day of the experiment all fry were collected to calculate the final length, weight and survival. 20 (twenty) fishes were randomly selected from each pond to determine the growth and survival.

2.6 Estimation of Length and Weight

The length gain, weight gain and daily weight gain were calculated as:

Length gain (cm) = (Mean final length – Mean initial length)

Weight gain (g) = (Mean final weight – Mean initial weight) and

Daily weight gain (g/day) = (Mean final weight – Mean initial weight) / Duration of the experiment (Number of the day).

Survival (%) = (Total no. of fry harvested/ Total no of fry released) X 100

2.7 Statistical Analysis

The mean values for the growth of different treatments were tested using one way ANOVA. Each value and parameters of different treatments expressed as mean \pm Standard deviation (SD). The level of significance was 95%.

3. RESULTS

3.1 Water Quality Parameters

There were no significant differences were observed in water quality parameters of different experimental ponds. Water quality parameters were within the best ranges during the sampling period (Table 2).

The growth and survival details of three treatments along with the initial and final particulars are presented in Table 3. In the experiment, *T. putitora* attained maximum length and weight in the lowest stocking density of 1.0×10^5 /ha in T₁ followed by T₂ and T₃ with the stocking densities of 1.5×10^5 and 2.0×10^5 /ha, respectively (Fig. 1 and Fig. 2). It was also observed higher daily weight gain in the lowest stocking density in T₁ followed by T₂ and T₃, respectively. There were significant (P<0.05) differences in final length of T₁, T₂ and T₃. In the case of final weight gain and daily weight gain, T₁ was significantly compared with T₂ and T₃; T₂ and T₃ also had significant differences. The highest survival rate (79.66 ± 4.34 %) also observed in the lowest stocking density (T₁) (Fig. 3). All treatments were implemented under the same environmental conditions and feeding was same. Ponds size and depth were also same for all treatments. Nursery management procedure was also the same for all the treatments. Only the stocking densities were different among the treatments and shows significant difference in final length, weight and daily weight gain (DWG).

All values were reported as mean and standard deviation ($M \pm SD$). Figures in the same row having the same superscripts are not significantly different and having different superscripts are significantly different ($P < 0.05$).

4. DISCUSSION

Growth in terms of length and weight was maximum at the lower stocking density and showed a significant decreasing trend with an increase in stocking density ($P < 0.05$). It is empirical that stocking density has a direct effect on food supply, space for living and water quality. Different studies indicate that there was a significant inverse linear relationship between stocking rate and growth linked such depression in growth to unavailability of proper space [13,14,6]. Physiological stress may also cause poor growth performance under high stocking densities as observed in *Catla catla* and *Labeo rohita* in cage culture [14]. Low water quality also causes poor growth in some cases of fry rearing [15,16]. In this experiment, the values of different water quality parameters were generally within the optimum requirements for the nursery rearing of Mahseer spawn [6]. In the present experiment, no distinct variations in the water quality parameters were observed at different stocking densities indicated that increasing stocking densities did not impose any environmental stress to the fish. There were the same findings also observed in *Carassius carassius* on growth and survival of primary nursing system [17]. However, the level of Dissolve oxygen (DO) was optimum in experimental ponds. Alkalinity level indicates the natural production of these ponds were moderately high. Which is closed to other carps fry rearing system [18,7]. Higher alkalinity values might be due to the higher amount of lime doses at the experimental period. The pH level of this experiment was optimum for the fish culture [6].

In this experiment, crude protein level having 32 % dry weights were used in supplementary feeding for *Tor putitora* spawn. For the growth of *Labeo rohita* spawn optimum protein requirement is 36 % [14]. 34.11% crude protein use in the one stage nursery management system of Gold fish [16]. Supplementary feeds containing 20.3-29.5% crude protein for a semi-intensive culture of *T. putitora* record the lower growth performance [18].

High growth performance was observed in terms of weight; length and weight gain of *Tor putitora* spawn in T_1 where the stocking density was lower as compared to T_2 and T_3 , although same food was supplied at an equal ratio in all the treatments due to the reason of competition for food at higher density treatments. T_1 having low density considering the suitable stocking density for the fish growth in nursery pond, as the number of spawn increases for the same stocking area, its effects the growth performance of Mahseer [6]. The treatments having great number of spawn with a high food concentration cause a stressful situation [7]. The highest weight gains of Mahseer (*Tor putitora*) fry and Crucian Carp (*Carassius carassius*) in nursery ponds in lowest the stocking density [4,17]. The highest survival in showed in lowest stocking density under single stage nursery system of Mahseer (*Tor putitora*) [7]. the same result observe in Crucian Carp (*Carassius carassius*) in primary nursery system [17].

The low density of *Tor putitora* fry that shows higher growth compare with higher density. Because higher density treatment fry have higher competition for food and space [13,19]. Further research should be done on stocking densities of this critically endangered species to understand the growth and survival in culture system of *T. putitora*.

Table 1. Proximate composition (% of dry matter) of the supplementary feed

Crude protein	Crude lipid	Crude fiber	Ash	NFE*
32.00	8.80	5.68	18.20	35.32

*NFE (Nitrogen free content) = $100 - \% (\text{crude protein} + \text{crude lipid} + \text{crude fiber} + \text{ash})$

Table 2. Water quality parameters of experimental ponds (31 December 2019- 10 March 2020)

Sl no.	Parameters	T_1	T_2	T^3
1	Temperature ($^{\circ}C$)	18.2- 31.7	18.0- 31.7	18.2- 31.6
2	Dissolve oxygen, DO (mg/l)	6.80- 7.45	6.80- 7.40	6.75- 7.40
3	pH	7.90- 8.64	7.90- 8.60	7.80- 8.60
4	Ammonia (mg/l)	0- 0.03	0- 0.03	0- 0.03
5	Alkalinity (mg/l)	160- 180	160- 180	160- 180




Fig. 1. Individual final length (cm) of *Tor putitora* fry under different stocking density treatments




Fig. 2. The final weight of *T. putitora* fry under different stocking density treatments

Table 3. Stocking, harvesting and growth performances of *T. putitora* in experimental ponds under different stocking densities

Treatment	Stocking density/ha	Released	Harvested	Length (cm)		Weight (g)		Daily weight gain (g/day)	Survival (%)
				Initial	Final	Initial	Final		
T ₁	1.0X10 ⁵	12,000	9,560	3.12± 0.65	8.67± 0.32 ^a	1.12± 0.12	5.74± 0.56 ^a	0.066± 0.006 ^a	79.66± 4.34 ^a
T ₂	1.5X10 ⁵	18,000	12,056	3.12± 0.65	7.89± 0.24 ^b	1.12± 0.12	4.42± 0.36 ^b	0.044± 0.004 ^b	66.97±3.67 ^b
T ₃	2.0X10 ⁵	24,000	13,123	3.12± 0.65	6.78± 0.25 ^c	1.12± 0.12	2.8± 0.37 ^b	0.024± 0.003 ^c	54.67± 3.12 ^c




Fig. 3. Survival (%) of *Tor putitora* fry under different stocking density treatments

5. CONCLUSIONS

From the above findings of this experiment, it can be concluded that the length and weight gain of *T. putitora* fry were inversely related to the stocking densities. Decreased stocking densities lead to an increase in length and weight. Finally, present findings of this experiment might be helpful for the fry production of *T. putitora* and take a little step to commercialization of its culture.

DISCLAIMER

The products used for this research are commonly and predominantly use products in area of research in Bangladesh. There is absolutely no conflict of interest between the authors and producers of the products because there are not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mohanty AN, Chatterjee DK, Giri BS. Effective combination of urea and

bleaching powder as a pesticide in aquaculture operation. *J. Aqua. Trop.* 1993;8:249-254.

2. Wedemeyer GA. Physiological response of juvenile Coho Salmon (*Oncorhynchus kisutch*) and Rainbow Trout (*Salmo gairdneri*) to handling and crowding stress in intensive fish culture. *J. Fish. Res. Board, Can.* 1976;33:2699-2702.
3. Rahman MA, Mazid MA, Rahman MR, Khan MN, Hossain MA, Hussain MG. Effect of stocking density on survival and growth of critically endangered mahseer, *Tor putitora* (Hamilton), in nursery pond. *Aquaculture.* 2005;249:275- 284.
4. Rahman MR, Rahman NA. Studies on the growth, survival and production of Calbasu (*Labeo calbasus* Ham.) fry at different stocking densities in primary nursing. *Bull. Fac. Sci., Univ. Ryuwus, Jpn.* 2003;76: 245- 255.
5. Ahmed N. Study on the spawning habits and early development of Copper Mahseer, *Barbus (Lissochilus) hexagonolepis*. McDonald. *Proc. National Institute of Science. India.* 1969;14:21-28.
6. Riar MGS, Raushon NA, Chowdhury P, Rahman MK. Effect of stocking density on growth performance and the survival of golden mahseer, *Tor putitora* (Hamilton) in primary nursing system. *International Journal of Fisheries and Aquatic Studies* 2019;7(5):200-203.

7. Rahman AKA. Freshwater Fishes of Bangladesh, 2nd ed., Zool. Soc. Bangladesh, Dhaka, Bangladesh. 2005; xviii:394.
8. Islam MS. Evaluation of supplementary feeds for semi-intensive pond culture of mahseer, *Tor putitora* (Hamilton). Aquaculture. 2002;212(1):263-276.
9. Hussain MG, Mazid MA. Genetic improvement and conservation of carp species in Bangladesh. Bangladesh Fisheries Research Institute and International Center for Living Aquatic Resources Management. 2001;74.
10. IUCN Bangladesh. Red List of Bangladesh Freshwater Fishes. IUCN, International Union for Conservation of Nature, Bangladesh Country Office, Dhaka, Bangladesh, 2015;5:xvi+360.
11. Kohinoor AHM. Development of culture technology of three small indigenous fish Mola (*Amblypharyngodon mola*), Punti (*Puntius sophore*) and Chela (*Chela cachius*) with notes on some aspects of their biology. PhD dissertation, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, Bangladesh. 2000;363.
12. AOAC (Association of Official Analytical Chemicals). Official Methods of Analysis of the Association of Official Analytical Chemists. 20th edition. Washington, DC; 2016.
13. Ronald N, Gladys B, Gasper E. The Effects of Stocking Density on the Growth and Survival of Nile Tilapia (*Oreochromis niloticus*) Fry at Son Fish Farm, Uganda. J Aquac Res Development. 2014;5:222. DOI: 10.4172/2155-9546.1000222
14. Mane AM, Dube K, Varghese T, Chavan BR, Kamble MT. Effects of Stocking Density on Growth Performance, Survival and Production of *Catla catla* and *Labeo rohita* During Nursery Rearing in Cages. Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci. 2019;89:275–281. Available:https://doi.org/10.1007/s40011-017-0940-2
15. Ferdous Z, Masum MA, Ali MM. Influence of Stocking Density on Growth Performance and Survival of Monosex Tilapia (*Oreochromis niloticus*) Fry. International Journal of Research in Fisheries and Aquaculture. 2014;4(2):99-103.
16. Shajib MSH, Sarker B, Asif AA, Rahman MM, Zafar MA, Hossain A. Effects of stocking density on the growth rate of gold fish fry reared in hapa. Asian J. Med. Biol. Res. 2017;3 (4):504-515.
17. Ali SM, Riar MGS, Pavel MSF, Mahmud Y. Primary nursery rearing of Crucian carp fry under different stocking densities. Journal Bangladesh Society of Agricultural Science and Technology. 2014;11(3&4):25-30.
18. Pathani SS. Studies on spawning ecology of Kumaun mahseer *Tor tor* (Hamilton) and *Tor putitora* (Hamilton). J. Bombay Nat. Hist. Soc. 1982;79:526- 530.
19. Chatta AM, Ahmad Z, Hayat S, Naqvi SA, Khan AM. Studies on survival of endangered indus Golden Mahseer (*Tor macrolepis*) and its impact on growth of other carps. The Journal of Animal & Plant Sciences. 2015;25(3):542-549.

© 2021 Riar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle4.com/review-history/73736>