



# Studies on Genetic Variability and Heritability in Rice Genotypes (*Oryza sativa*. L.) under Salinity

P. Hima Bindu <sup>a\*</sup>, R.M. Sundaram <sup>b</sup>, G. Shiva Prasad <sup>c</sup>,  
C.H. Damodar Raju <sup>d</sup> and K. Sumalini <sup>a</sup>

<sup>a</sup> Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad – 500030, India.

<sup>b</sup> ICAR-Indian Institute of Rice Research, Rajendranagar, Hyderabad -500030, India.

<sup>c</sup> Agricultural Research Station, Kampasagar, Nalgonda -508001, India.

<sup>d</sup> Rice Breeding, Regional Agricultural Research Station, Warangal -506007, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101524>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124944>

Original Research Article

Received: 06/08/2024

Accepted: 08/10/2024

Published: 15/10/2024

## ABSTRACT

Rice (*Oryza sativa* L.) is one of the most important staple food crops in the world because more than half of the world's population depends on rice for their livelihood. By 2050, the world's rice productivity will need to double in order to keep up with population growth. Narrow genetic base in the released variety has made the improvement in plateaus. In order to go more than the yield barrier, the genetic base must be broadened. Heritability and genetic variability were measured in eighty-six rice genotypes in terms of yield and yield contributing traits under saline soils. Significant differences were found between the genotypes for all of the studied traits according to analysis of

\*Corresponding author: E-mail: [himabindu4agri@gmail.com](mailto:himabindu4agri@gmail.com);

Cite as: Bindu, P. Hima, R.M. Sundaram, G. Shiva Prasad, C.H. Damodar Raju, and K. Sumalini. 2024. "Studies on Genetic Variability and Heritability in Rice Genotypes (*Oryza Sativa*. L.) under Salinity". *Journal of Advances in Biology & Biotechnology* 27 (10):1007-16. <https://doi.org/10.9734/jabb/2024/v27i101524>.

variance. High PCV and GCV were recorded for the traits viz., mortality percentage (%), total number of grains per panicle, number of un-filled grains per panicle, sterility percentage (%) and seed yield per plant (g). The traits showed a significant degree of genetic advancement together with high heritability were mortality percentage (%), 1000 grain weight (g) and seed yield per plant (g). Genetic advance over mean were high for mortality percentage (%), number of productive tillers/m<sup>2</sup>, total number of grains per panicle (g), number of un-filled grains per panicle (g), sterility percentage (%), 1000 grain weight (g) and seed yield per plant (g). This study provides information about the variability and heritability measures which may supplement the breeding program to break the yield plateaus.

**Keywords:** Genetic variability; genetic advance; heritability; salinity; rice.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is the primary food source for one-third of the world's population and 70% of Indians, India was determined to have 6.7 million hectares damaged by salt. Of which 3.7 mha were sodic and 2.9 mha were saline [1]. 43.79 million hectares of land are utilized to grow rice in India, with a yield of 116.42 million tonnes and 2.7 tonnes per hectare. Salt stress severely affects crop development by lowering photosynthetic potential of plants. High sodium content in soil damages cell metabolism and photosynthesis by increasing the toxicity of sodium in the cytosol and producing osmotic stress on cell-water relations [2]. Na<sup>+</sup> and K<sup>+</sup> exhibit comparable ionic, physical, and chemical characteristics. K<sup>+</sup> plays key part in numerous physiological process and cell metabolism for optimal growth and development. When roots are under salt stress, their Na<sup>+</sup> concentration goes over what is ideal, which lowers their ability to absorb K<sup>+</sup>. As a result, various plant portions retain less K<sup>+</sup>. Toxic effects result from Na<sup>+</sup> entering the K<sup>+</sup> route and changing the ion ratio in the cytosol. Extrusion of surplus Na<sup>+</sup> through roots and compartmentalization in distinct regions are two mechanisms used by salt-tolerant plants to maintain low cytosolic Na<sup>+</sup> / K<sup>+</sup> ratios, which in turn lowers Na<sup>+</sup> concentration in the cytosol. By overcoming the negative effects of salt, this osmotic adjustment helps to provide a significant yield [3]. Local cultivar variability is declining as a result of the ongoing cultivation of high-yielding cultivars.

Plant breeding is based on variation. Any crop improvement program's ability to succeed is mostly determined by the amount and range of genetic stock variability. In crop improvement programs, the initiation of suitable breeding processes requires a crucial estimation of genetic diversity. In order to help breeders plan

an appropriate breeding program, it becomes necessary to split overall variability into its heritable and non-heritable components using certain genetic criteria. As a result, the degree and type of genetic variability within a population determines its rate of development. Information of genetic variability among growth as well as yield components in rice has been reported by many workers [4].

The degree of variation in the population can be determined using genetic variability parameters including Heritability, Phenotypic Coefficient of Variation (PCV), Genotypic Coefficient of Variation (GCV), and Genetic Advance as percent Mean (GAM). Plant breeders have not yet completely utilized the rich pool of useful genes that have collected throughout time in the rice germplasm to improve varieties. In order to support future breeding programs for yield enhancement through expanding the genetic base and genetic recombination, the goals of this study were to evaluate genetic variability, heritability, and genetic advancement.

## 2. MATERIALS AND METHODS

During *Rabi*, 2020-21, 86 rice genotypes along with checks were sown in the nursery bed at Agricultural Research Station, Kampsagar (Table 1). Twenty-one day old seedlings were transplanted in a Randomized Block Design, under saline field conditions (pH: 9.30, E.C: 4.68 dSm<sup>-1</sup>, and ESP: 88.0) with a spacing of 20 x 15 cm. General view of field screening was shown in plate 1. The crop was grown according to the recommended set of cultural practices. During reproductive stage, visual scoring of salt injury was also recorded according to IRRI-SES 2013 (Table 2). Observations on yield and yield attributing characters were recorded while screening for salinity tolerance at the reproductive stage.

**Table 1. List of rice genotypes and checks utilized in the screening experiment**

S.No	Genotype	S.No	Genotype	S.No	Genotype
1.	IR69726-116-1-1	30.	KPS10329	59.	IR 10F 360
2.	IR77186-122-2-2-3	31.	MINGHU163	60.	DJ 123
3.	NSICRC240	32.	ZHENSHAN 97B	61.	Oryzica 1
4.	IRRI154	33.	IR64-21	62.	FEDEARROZ 50
5.	GSRIR2-9-RI-SU3-Y2	34.	IRBB 66	63.	TEQING
6.	CT118911-2-2-7-M	35.	IR 78222-20-7-148-2-B-B-B-B	64.	UPL RI7:IRTP9897-C1
7.	IR13F167	36.	IRRI 147	65.	Jamir
8.	Sahel177	37.	SANHUANGZHAN NO2	66.	IRIOM300
9.	Jasmine85	38.	IR77298	67.	KPS 13575
10.	M202	39.	Sambha mahsuri+sub1	68.	KPS 13576
11.	KPS10628	40.	SUPA	69.	KPS 13577
12.	KPS10631	41.	IRRI 104	70.	KPS 13578
13.	KPS10633	42.	N 22 IRGC 19379-1	71.	KPS 13579
14.	KPS10640	43.	MTU 1010	72.	KPS 13580
15.	KPS10642	44.	SWARNA	73.	KPS 13581
16.	KPS10651	45.	NANHI	74.	KPS 13582
17.	KPS10654	46.	Kinandang patong	75.	KPS 13583
18.	KPS10656	47.	SADRI	76.	KPS 13584
19.	KPS10657	48.	OM4900	77.	KPS 13585
20.	KPS10658	49.	IR93340	78.	KPS 13586
21.	KPS10661	50.	IR93354	79.	KPS 13587
22.	KPS10667	51.	Khao Hlan ON	80.	KPS 13588
23.	KPS10669	52.	IR84984-83-15-481-B		<b>CHECKS</b>
24.	KPS10672	53.	Manaw Thuka	1.	FL478
25.	KPS10676	54.	BR 28	2.	PUSA 44
26.	KPS10683	55.	TN1	3.	CSR 23
27.	KPS10316	56.	IR 6	4.	CSR 36
28.	KPS10319	57.	Zanton:IRGC31248-1	5.	RNR11718
29.	KPS10321	58.	URAIBOOL IRGC52785-1	6.	KPS2874

**Table 2. Scoring of damage for salt injury in field conditions in rice standard evaluation system scale (IRRI-SES 2013)**

Score	Growth Scale	Salinity-induced reaction
1	Normal growth, no leaf symptoms	Highly tolerant
3	Nearly normal growth, but leaf tips of few leaves whitish and rolled.	Tolerant
5	Growth severely retarded, most leaves rolled, only a few are elongating	Moderately tolerant
7	Complete cessation of growth, most leaves dry, some plants drying	Susceptible
9	Almost all plants dead or drying	Highly susceptible

Data were collected on the following: percentage of mortality (M%), days to 50% flowering (DFF), plant height (PH), length of panicle (PL), number of productive tillers (NPT), number of grains per panicle (NGP), number of unfilled grains per panicle (UFG), spikelet sterility (S%), 1000 grain weight (TW), and seed yield per plant (SY). Each entry consisted of five randomly selected plants for observations, from which the mean value was calculated.

Analysis of variance and estimates of the phenotypic and genotypic coefficient of variation were estimated using the techniques suggested by Burton [5] and Panse and Sukhatme [6]. For the features under study, estimates of heritability in the wide sense (Lush, 1940) and genetic advancement as a percentage of mean [7] were also made. According to Johnson et al., [7] the range of heredity was categorized as low (10–30%), medium (30–60%), and high (>60%).

According to Johnson et al., [7] the genetic progress for the traits was categorized as strong (>20%), moderate (10-20%), or low (<10%). The software INDOSTAT version 9.2 was used to do the statistical analysis.

### 3. RESULTS AN DISCUSSION

For the success of any breeding programme, it is essential to know the inherent genetic variability in a given crop species for a particular trait under development is essential. The coefficient of variation is used to estimate the degree of variation existing in the crop species. Estimates of heritability provide information on the amount of transmissible genetic variation among total variation, which defines its responsiveness to selection and the scope of improvement in the new population over the original population through genetic advance. Analysis of variance

showed that the mean sum of squares for genotypes were highly significant for all the characters studied (Table 3). It indicates that presence of high variability for the traits in rice genotypes used in the study.

If a character is driven by non-additive gene action, it may have high heritability but low genetic advance, but if it is governed by additive gene action, it will have high heritability (above 60%) and high genetic advance (above 20%), provides much potential for further improvement. As a result, the most significant strategy to make genetic gain through generations is to increase these qualities through selection. For the genotypes, the genotypic and phenotypic coefficients of variation, heritability, genetic advance, and genetic advance as a percentage of the mean were calculated and shown (Table 4; Figs. 1 &2).



Plate 1. Over view of salinity block





Fig. 1. Genotypic and phenotypic coefficients of variation for yield and yield contributing traits in rice genotypes



**Fig. 2. Heritability and genetic as per cent of mean (5%) for yield and yield attributing traits in rice genotypes**

The GCV and PCV values for mortality percentage were high at 30.52% and 33.97% respectively. Heritability values for this trait were high at 80.71% with a moderate genetic advance of 15.20% and a high genetic advance as a per cent of mean 56.47%. As this characteristic has a negative impact on yield, it should be subjected to negative selection pressure in order to develop varieties with high yield tolerant to salinity stress.

The GCV and PCV values for days to 50% flowering were both low in magnitude, at 1.92% and 5.02%, respectively. This wide range of variation provides shifting the seasonal timing of reproduction and produce novel varieties that are better adopted to local environment and change in climatic conditions. This trait had a low heritability of 14.70%, with a low genetic advance of 1.53% and a poor genetic advance as a per cent of the mean of 1.52%. As heritability and genetic advance are low, it indicates the preponderance of non-additive gene action, therefore selection for this trait is inefficient due to non-additive gene action. These results are in confirmity with the findings of Parimala and Devi [8] and Hasan-Ud-Daulla and Sarker [9].

“The GCV and PCV values for plant height were 6.97% and 9.09%, respectively, which were low in magnitude. The trait's heritability value was moderate 58.82%, with moderate genetic advance 10.23% and genetic advance as a percentage of the mean 11.01%. When

heritability is moderate and GAM is moderate, it implies that the environment has control over the character, and that improvement can be achieved by intermitting superior genotypes of segregating populations developed through combination breeding. These results are in trend with the findings” of Singh and Verma [10] and Saha et al. [11].

“The GCV and PCV values for the trait panicle length were 7.30% and 13.34% with low and moderate magnitude respectively. This trait heritability was moderate 30.01%, with low genetic advancement 1.68% and genetic advance as per cent of mean 8.23% respectively. Moderate values of heritability and low genetic advance indicate a lack of inherent variability and limited scope for improvement through selection for these traits among the lines”. Sumanth et al. [12] and Divya et al. [13] also obtained similar results.

“The GCV and PCV values for the trait number of productive tillers/m<sup>2</sup> were 14.94% and 19.89%, respectively, which were moderate in magnitude. This trait had a moderate heritability of 56.40%, and high genetic advance of 39.36%, with high genetic advance as a per cent of mean of 23.11%. The predominance of non-additive gene action is shown by moderate estimates of heritability and high genetic advance. This trait can be improved by heterosis breeding. These results are in confirmity with those” of Parimala and Devi [8] and Swapnil et al. [14].

**Table 3. Analysis of variance for yield and its attributing traits in rice genotypes evaluated under saline field conditions**

	<b>DF</b>	<b>M</b>	<b>DFF</b>	<b>PH</b>	<b>NPT/m<sup>2</sup></b>	<b>PL</b>	<b>NGP</b>	<b>UFG</b>	<b>S</b>	<b>TW</b>	<b>SY</b>
Replicates	2	47.16	8.49	37.68	360.81	100.35*	23.49	75.76	40.39	26.79	31.24
Treatments	85	218.48**	33.17*	156.93**	2443.44**	11.78**	358.34**	176.40**	456.52**	30.44**	41.53**
Error (A)	170	16.13	21.88	29.74	500.83	5.16	81.57	56.12	154.10	2.23	7.47
Total	257	83.30	25.51	71.87	1142.24	8.09	172.66	96.05	253.23	11.75	18.92
General Mean		26.91	100.95	93.39	170.35	20.34	62.67	20.51	33.49	19.89	13.11

\* - significant at 5 per cent level, \*\* - significant at 1 per cent level DF- Degrees of freedom

M (%) - Mortality percentage

DFF- Days to 50 % flowering

PH (cm) - Plant height

PL (cm) - Panicle length

NPT/ m<sup>2</sup> - Number of Productive tillers/m<sup>2</sup>

NGP - Total number of grains/panicle

UFG - Number of un-filled grains/panicle

S (%) - Sterility percentage

TW (g) - 1000 grain weight

SY (g) - Seed yield/plant

**Table 4. Estimates of genetic parameters for yield and yield attributing traits in rice genotypes evaluated under saline field condition**

S.No.	Traits	PCV (%)	GCV (%)	h <sup>2</sup> (%)	GAP (%)	GAM (%)
1.	Mortality percentage (%)	33.97	30.52	80.71	15.20	56.47
2.	Days to 50 % flowering	5.02	1.92	14.70	1.53	1.52
3.	Plant height (cm)	9.09	6.97	58.82	10.23	11.01
4.	Panicle length (cm)	13.34	7.30	30.01	1.68	8.23
5.	Number of Productive tillers/m <sup>2</sup>	19.89	14.94	56.40	39.36	23.11
6.	Total number of grains/panicle	21.04	15.33	53.12	14.42	23.00
7.	Number of un-filled grains/panicle	47.82	30.87	41.74	8.42	41.05
8.	Sterility percentage (%)	47.67	29.98	39.51	13.00	38.84
9.	1000 grain weight (g)	17.14	15.41	80.80	5.68	28.54
10.	Seed yield/plant (g)	33.08	25.69	60.31	5.40	41.06

PCV= Phenotypic coefficient of variation, GCV= Genotypic coefficient of variation, h<sup>2</sup>=Broad-senseHeritability, GAP= Genetic advance per cent (at 5%), GAM= Genetic advance as a per cent of the mean (at 5 %)

“The GCV and PCV values for the trait number of grains per panicle were 15.33% and 21.04% respectively which were moderate and high in magnitude respectively. Heritability values for this trait were moderate at 53.12% and with a moderate genetic advance of 14.42% and high genetic advance as per cent of mean 23.00%. Moderate heritability and high genetic advance for this character indicate predominance of non-additive gene action in the control of this character it can be improved by heterosis breeding. Similar kind of results were reported” by Sameera et al. [15] and Saha et al. [11].

“The GCV and PCV values for the trait un-filled grains per panicle were 30.87% and 47.82%, respectively, which were high. This trait's heritability was moderate 41.74%, with a moderate genetic advance of 8.42% and a high genetic advance as a per cent of the mean 41.05%, indicating a good response to selection”. Hasan-Ud-Daula and Sarker [9] obtained similar results.

“The GCV and PCV values for trait sterility percentage were both high in magnitude, at 29.98% and 47.67% correspondingly. Negative selection can be utilised efficiently with heritability values of moderate 39.51%, and moderate genetic advance of 13.00%, and high genetic advance as a per cent of mean of 38.84%. As this trait has a negative impact on yield, it should be subjected to negative selection pressure in order to boost develop lines with sterility tolerance to yield”. Hasan-Ud-Daula and Sarker [9] obtained similar results.

“The GCV and PCV values for the 1000 grain weight were 15.41% and 17.14% respectively which were moderate in magnitude. Heritability values for this trait were high 80.80% with a low genetic advance of 5.68% and high genetic advance as per cent of mean 28.54%. Results indicated possible scope for improvement through selection of this trait and breeder may have reliable benefits in next generation with respect to this trait. These findings were in accordance with those” of Rashid et al. [16] and Sumanth et al. [12].

“The GCV and PCV values for the trait seed yield per plant were 25.69% and 33.08% respectively, both were quite high. This characteristic had a high heritability of 60.31%, with a low genetic advance of 5.40% and a high genetic advance as a per cent of the mean of 41.06%. Yield, as a complex character, was influenced by its attributing factors and was shown to have high heritability and GA, indicating that it is controlled by additive genetic effects and a simple phenotypic selection will hasten the varietal improvement period. Similar results were recorded” by Bhati et al. [17] and Tiwari [18].

The PCV values in this study were higher or close to the corresponding GCV values for all characteristics. The small differences between PCV and GCV for all the traits indicated that the variability was predominately due to genotypic differences, *i.e.*, fewer environmental influences. In the present investigation, high values of GCV and PCV were observed in the case of mortality percentage (%), total number of grains per panicle, number of un-filled grains per panicle,

sterility percentage (%) and seed yield per plant (g) indicating the presence of inherent variability among the lines. The difference between PCV and GCV was high for un-filled grains and sterility percentage (%), indicating the influence of environment on these traits *i.e.*, indicating the presence of saline environment effects on the treatments under study. These findings of high values of GCV and PCV for un-filled grains per panicle, and sterility percentage (%) were in accordance with the findings of Hasan-Ud-Daula and Sarker [9]. Moderate estimates of GCV and PCV were observed in the case of Sumanth et al. [12] for 1000 grain weight, Parimala and Devi [8] for number of productive tillers/m<sup>2</sup>. Low estimates of GCV and PCV were observed in the case of days to 50 % flowering, plant height (cm), and panicle length (cm), indicating a lack of inherent variability and limited scope of improvement through selection for these traits among the genotypes. Similar results were observed by Parimala and Devi [8] and Hasan-Ud-Daula and Sarker [9] for days to 50 % flowering, Singh and Varma [10] Saha et al. [11] for plant height, Sumanth et al. [12] and Divya et al. [13] for panicle length.

Higher estimates of heritability for characters such as mortality percentage, 1000 grain weight (g) and seed yield per plant (g) indicated that these characters are less influenced by environmental fluctuations. Hence, selection for improvement of such characters may not be useful as broad sense heritability depends on total genetic variance which includes both additive and non-additive variances. Senguttuvel et al. [19] Akinwale et al. [20] and Hasan-Ud-Daula and Sarker [9] reported the same results for heritability for seed yield per plant.

An estimated heritability values alone is less reliable as these values are prone to alter with change in the environment and experimental material. Hence, the use of high heritability coupled with high genetic advance as per cent of mean is preferred and most probably controlled by additive gene action which may respond to selection.

Genetic advance was low for days to 50% flowering, panicle length (cm), number of un-filled grains per panicle, 1000 grain weight (g) and seed yield per plant (g). Moderate values were observed in the case of traits such as mortality percentage (%), plant height (cm), total number of filled grains per panicle, and sterility percentage (%). A high value of genetic advance

was observed in the case of number of productive tillers/m<sup>2</sup>. A similar case of the low estimate for days to 50 % flowering was reported by Hasan-Ud-Daula and Sarker [9] whereas, Divya et al. [13] and Singh and Verma [10] reported it in case of plant height, Manohara and Singh [21] for the total number of grains per panicle and Parimala and Devi [8] for seed yield per plant (g).

“The estimates of genetic advance over mean were high for mortality percentage (%), number of productive tillers/m<sup>2</sup>, total number of grains per panicle (g), number of un-filled grains per panicle (g), sterility percentage (%), 1000 grain weight (g) and seed yield per plant (g), indicating that these characters are governed by non-additive gene action and heterosis breeding programme may be useful to improve these traits. These findings align with the reports” of Sumanth et al. [12] Devi et al. (2017), Islam et al. (2016), Divya et al. [13] Hasan-Ud-Daula and Sarker [9].

Characters having a high heritability and a high genetic advance over the mean are more reliable for selection. The traits mortality percentage (%), 1000 grain weight (g) and seed yield per plant (g) were observed to be having high heritability coupled with high genetic advance. Partitioning variability into fixable and non-fixable components is critical for understanding the true picture of trait genetic variation. High heritability with moderate GAM may be attributable to additive gene action and manifested as a result of the environment's favourable effect rather than genotype.

#### 4. CONCLUSION

The current investigation concluded that maximum values of GCV and PCV for mortality percentage, number of un-filled grains per panicle, sterility percentage and seed yield/plant denotes the occurrence of large amount of variation amongst the germplasm lines for potential yield enhancement through selection and also grain yield/plant perceived maximum heritability combined with high genetic advance as percent of mean, which revealed that this trait is regulated through additive gene action and thus, the selection might be practiced based on phenotypic performance.

Thus, it was found that traits possessing both strong genetic advancement and high heritability are controlled by additive gene action, which might be improved by applying simple selection



techniques. The presence of non-additive gene action is indicated by characteristics exhibiting high heritability with modest genetic advance. High environmental impacts have also revealed high heritability, thus selection for these traits must be successful.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Mandal A, Obi Reddy G, Ravisankar T. Digital database of salt affected soils in India using Geographic Information System. *Journal of Soil Salinity and Water Quality*. 2011;3(1):16-29.
2. Gupta P, Verma O, Verma R, Gupta R, Singh V, Jyoti K, Yadav R. Heritability and genetic advance analysis using generation mean analysis in rice (*Oryza sativa* L.) under sodic soil. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(5):1471-1475.
3. Chakraborty K, Basak N, Bhaduri D, Ray S, Vijayan J, Chattopadhyay K, Sarkar RK. Ionic basis of salt tolerance in plants: nutrient homeostasis and oxidative stress tolerance. *Plant nutrients and abiotic stress tolerance*. 2018;325-362.
4. Sivasubramanian S, Madhavamenon P. Combining ability in rice. *Madras Agricultural Journal*. 1973;60:419-421.
5. Burton GW. Quantitative inheritance in grasses. *Proceeding on 6th International Grassland Congress Journal*. 1952;1:277-283.
6. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. ICAR, New Delhi. 1985;236-246.
7. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybeans. *Agronomy Journal*. 1955;47(7):314-318.
8. Parimala K, Devi KR. Estimation of variability and genetic parameters in indica and japonica genotypes of rice (*Oryza sativa* L.). *International Journal of Current Microbiology and Applied Sciences*. 2019; 8(3):1138–1142.
9. Hasan-Ud-Daula H, Sarker U. Variability, heritability, character association, and path coefficient analysis in advanced breeding lines of rice (*Oryza sativa* L.). *Genetika*. 2020;52(2):711-726.
10. Singh N, Verma OP. Genetic variability, heritability and genetic advance in rice (*Oryza sativa* L.) under salt stressed soil. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(3):3114-3117.
11. Saha S, Hassan L, Haque M, Islam M, Rasel M. Genetic variability, heritability, correlation and path analyses of yield components in traditional rice (*Oryza sativa* L.) landraces. *Journal of the Bangladesh Agricultural University*. 2019; 17(1):26-32.
12. Sumanth V, Suresh BG, Jalandhar BR, Srujana G. Estimation of genetic variability, heritability and genetic advance for grain yield components in rice (*Oryza sativa* L.) *Journal of Pharmacognosy and Phytochemistry*. 2017;6(4):1437-1439.
13. Divya J, Raju Ch.D, Subba Rao LV, Aparna K. Genetic variability studies in genetically diverse rice genotypes. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(4):1169–1172.
14. Swapnil, Prasad K, Chakraborty M, Singh D, Kumari P, Ekka J. Genetic variability, correlation and path coefficient studies in F<sub>2</sub> generation of rice (*Oryza sativa* L.). *International Journal of Chemical Studies*. 2020;8(4):3116-3120.
15. Sameera SK, Srinivas T, Rajesh AP, Jayalakshmi V, Nirmala PJ. Variability and path co-efficient for yield and yield components in rice. *Bangladesh Journal of Agricultural Research*. 2016;41(2):259–271.
16. Rashid MM, Hassan L, Begum SN. Phenotypic performance of rice landraces under salinity stress in reproductive stage. *Progress Agriculture*. 2017;28(1):1-6.
17. Bhati M, Babu GS, Aishwarya SR. Genetic variability, correlation and path coefficient for grain yield and quantitative traits of elite rice (*Oryza sativa* L.) genotypes at Uttar Pradesh. *Electronic Journal of Plant Breeding*. 2015;6(2):586- 591.
18. Tiwari JK. Association analysis and selection strategies for various yield

- contributing traits in rice genotypes. Applied Biological Research. 2017;19(1): 35-40.
19. Senguttuvel P, Raveendran M, Vijayalakshmi C, Thiyagarajan K, Bapu JRK, Viraktamath BC. Molecular mechanism of salt tolerance for genetic diversity analysed in association with Na<sup>+</sup>/ K<sup>+</sup> ratio through SSR markers in rice (*Oryza sativa* L.). International Journal of Agricultural Research. 2010;5(9):708-719.
  20. Akinwale MG, Gregorio G, Nwilene F, Akinyele BO, Ogunbayo SA, Odiyi AC. Heritability and correlation coefficient analysis for yield and its components in rice (*Oryza sativa* L.). African Journal of Plant Science. 2011;5(3):207-212.
  21. Manohara KK, Singh NP. Genetic variability, correlation and path analysis in rice (*Oryza sativa* L.) under coastal salinity conditions of Goa. Journal of Indian Society for Coastal Agricultural Research. 2015;33(1):34–39.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/124944>