



Phytochemical Evaluation and *In-silico* Studies of Bioactive Compound from the Seeds of *Afrostryrax lepidophyllus* to Identify Potential Anti-inflammatory Agents

Ikezu, Ujupaul, Joy, M. ^{a*} and Ikpa Chinyere, Benardette, C. ^a

^a Department of Chemistry, Imo State University Owerri, Imo State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118460>

Original Research Article

Received: 07/04/2024

Accepted: 11/06/2024

Published: 15/06/2024

ABSTRACT

Afrostryrax lepidophyllus commonly known as country onions from the *Huaceae* family has been acclaimed to be used in treatment of several health conditions such as inflammation, prevention of infection and cancer, improve digestion, use traditionally as remedy for pain relief etc ethnomedically. This study was carried out to determine the bioactive composition of *A. lepidophyllus* seed and to evaluate the anti-inflammatory activities of seed extract using GC-MS) and in silico molecular docking. The GC-MS result revealed the presence of 18 bio-active compounds of which eugenol (47.14%), 3-ally-6-methoxyphenol (24.23%), Caryophyllene (10.35%) and oleic acid (7.91%) were the most abundant. The result revealed that the efficacy of the plant against inflammatory diseases could be linked to eugenol, 3-ally-6-methoxyphenol and Caryophyllene found in the crude extract. The bioactive compounds from the extract were docked

*Corresponding author: Email: joymag16@gmail.com;

Cite as: M. , Ikezu, Ujupaul, Joy, and Ikpa Chinyere, Benardette, C. 2024. "Phytochemical Evaluation and In-Silico Studies of Bioactive Compound from the Seeds of *Afrostryrax Lepidophyllus* to Identify Potential Anti-Inflammatory Agents". *Journal of Materials Science Research and Reviews* 7 (2):238-51. <https://journaljmsrr.com/index.php/JMSRR/article/view/328>.

on the inflammatory protein crystal structure of diclofenac bound to the cyclooxygenase active site of COX-2 (1pxx) to know their binding affinities and compared with that of some known anti-inflammatory drugs. The docking result showed that the standard drugs Diclofenac has the best binding affinity of -8.5 kcal/mol, followed by Naproxen (-8.1kcal/mol), Caryophyllene (-7.5kcal/mol), Bicyclo [7.2.0] undec-4-ene (-7.4 kcal/mol) have better affinity than the standard drug Ibuprofen (-7.3 kcal/mol). All the compounds from the exact have high binding affinities within the range of -7.5 – 6.1kcal/mol except 2,4,5,7-tetrathiaoctane that is -3.5 kcal/mol. The interaction results of the extracts with the diseased protein proved the anti-inflammatory efficacy of the seed as acclaimed therefore validating the traditional use of the seed against inflammatory diseases.

Keywords: Spices; bioactive compounds; anti-inflammatory; docking; interaction.

1. INTRODUCTION

Spices and herbs have been an essential part in human life for centuries both for culinary and medicinal purposes, they have been used at a domestic and industrial level as flavoring, preservation, and coloring agent in nutraceutical, pharmaceutical, and cosmetics products [1]. Spices not only enhance the flavor, aroma, and color of food and beverages, but they can also give protection from acute and chronic diseases. More Americans are considering the use of spices and herbs for medicinal and therapeutic/remedy use, especially for various chronic conditions. There is now ample evidence that spices and herbs possess antioxidant, anti-inflammatory, antitumorigenic, anticarcinogenic, and glucose and cholesterol lowering activities as well as properties that affect cognition and mood [2,3-5]. Research over the past decade has reported on the diverse range of health properties possessed by herbs and spices via their bioactive constituents, including sulfur-containing compounds, tannins, alkaloids, phenolic diterpenoids, and vitamins, especially flavonoids and polyphenols. Spices and herbs such as clove, rosemary, sage, oregano, and cinnamon are excellent sources of antioxidants with their high content of phenolic compounds [6]. It is evident that frequent consumption of spicy foods was also linked to a lower risk of death from cancer and ischemic heart and respiratory system diseases [7]. Among these spices is the Country Onions, scientifically known as *Afrostryax lepidophyllus* from *Huaceae* family, a culinary treasure from the vibrant rainforests of Central and West Africa which is perfect for adding depth and richness to a variety of dishes. This onion stands out with its unique garlic-like flavor profile, infusing a distinctive pungency into traditional dishes. It is oval in shape with a brown bark, the seed is widely used as a flavoring ingredient in many traditional dishes because of its strong and pungent aroma. Country Onion is a bulbous plant that belongs to the *Allium* family,

which also includes garlic, onions, and chives. It is native to Africa where it is an important traditional food source for local communities, Country onions is a flavoural and nutritive ingredient, widely used in different parts of Africa, including Nigeria, Cameroon and, Ghana for centuries as a staple ingredient in their cuisine. People love this charming spice for their unique flavors and aroma, which resemble garlic more than traditional onions. But what truly sparks curiosity is their hidden health benefits [8]. They have a long history as cherished ingredients in African cuisines and, beyond their culinary charm, offer notable health benefits.

Country onions has been reported as a rich source of antioxidants containing compounds like allicin and alliin, which have been shown to have antibacterial, antiviral, and antifungal properties. Studies have also suggested that Country Onion can help lower blood pressure and cholesterol, reduce inflammation, boost immune system, possessed anticancer properties and improve cardiovascular health. It is available in Cameroon, Gabon, Nigeria and Ghana. Country onions is currently listed by the International Union for Conservation of Nature as “vulnerable”, giving it high conservation value [9]. *Afrostryax lepidophyllus* is also known as Olum or Bombimbi. This work is aimed at identifying the bioactive component of this treasured spice. Insilico methods was used to study the anti-inflammatory potentials of the plant phytocompounds and other commercial drug to ascertain its efficacy as acclaimed ethnomedically and by previous researchers.

2. MATERIALS AND METHODS

Matured seeds of *Country Onion* were bought from a local market in Anambra State. The dried seeds were subjected to preliminary grinding using mechanical hand grinder, and the particle size were reduced further using an electric

blender. The pulverized seed were stored in an airtight plastic container for further analysis.

2.1 Extraction of Phytochemicals

The powdered *Country Onion* 200g was measured and percolated in a stoppered container containing 500ml of redistilled ethanol (98%) and allowed to stand at room temperature for a period of 3 days with frequent agitation until the soluble matter had dissolved. The mixture was then clarified by filtration and later concentrated with rotary evaporator at 55°C to get the crude sample for GCMS analysis [10].

2.2 GC-MS Analysis

The GC-MS analysis was done at Zaria, kaduna state Nigeria. The compounds in the sample were identified using agilent GC-MS (Agilent 19091-433HP, USA) coupled to a mass spectrophotometer. The initial column temperature was 35°C with a hold time of 3 minutes. The temperature was programmed to rise by 8°C/min with a final temperature of 280°C. In the process, 1µl of the sample was injected into the port and immediately vaporized and moved down the column with helium as the carrier gas with flow rate of 1 ml/min. The MS Spectrum was taken at 70 eV. The identification of the compounds was done by comparing the spectrum of unknown compounds with the spectrum of known compounds in NIST structural library [11].

The compounds identified from GCMS and their structures are shown in Table 1.

2.3 Identification and Preparation of Ligands

The 3D structure-data files (SDF) of the compounds in the crude extract sample and some anti-inflammatory drugs were identified and downloaded from the PubChem database. They were minimized in PyRx virtual screening tool, using Universal Force Field at 200 steps and converted to AutoDock ligands (pdbqt) and then used for the docking analysis. Identification and preparation of molecular targets crystal structures of diclofenac bound to the cyclooxygenase active site of cox-2 with protein ID:1pxx was identified and downloaded from the Protein Data Bank (PDB). The interfering crystallographic water molecules and cocrystallized ligand were removed, and minimization of the energy of the protein was then done using Biovia Discovery studies [12].

2.4 Docking Procedure and Analysis of Results

The screening of the phytochemical compounds from the seed extract was performed by docking them on selected binding pockets of proteins of crystal structures of diclofenac bound to the cyclooxygenase active site of cox-2 and ranked based on their binding energies. The multiple docking of the ligands and proteins was done with Autodock Vina in PyRx software. A rigid-flexible docking was performed after setting a grid box surrounding the binding sites of the receptors at exhaustiveness = 8, center x = 16.72, center y = 50.18, center z = 67.80, size x = 22.42, size y = 21.10, size z = 22.41. The molecular docking results were organized on an Excel spreadsheet, and the Heat Map of the data and the interaction was viewed using the Biovia discovery studio.

2.5 The Protein

The protein used for the docking is a high-resolution X-ray crystal structures of diclofenac bound to the cyclooxygenase active site of cox-2 (1pxx) officially known as prostaglandin-endoperoxide synthase (PTGS), an inflammatory protein. Inflammation is a helpful recovery process that cells utilize to stop the progression of harm or injury to tissues caused by foreign invaders and start the healing process. It's a complicated process involving white blood cells, macrophages, and inflammatory cytokines such as prostaglandins, TNF- (Tumor Necrosis Factor), interleukin IL6, and IL-8, to mention a few. The mobilization of arachidonic acid for prostaglandin production is a hallmark of inflammation. Cyclooxygenases are recognised to be the primary mediators of prostaglandin production, which are inflammatory indicators and are hence the focus of anti-inflammatory therapy. Cyclooxygenases such as COX-1 and COX-2 enzymes convert arachidonic acid to prostaglandins. COX-1 is required for the body's homeostatic activities, such as platelet synthesis for blood, kidney development and function, gastric mucosa maintenance, and so on. Increased inflammation, angiogenesis, metastatic and proliferative invasion, decreased apoptosis, and the establishment of an immunosuppressive microenvironment are all linked to COX-2 derived prostaglandin PGE2. A variety of drugs inhibit the conversion of arachidonic acid to prostaglandin G2 by the cyclooxygenase (COX) activity of prostaglandin endoperoxide synthases.




Plate 1. Unprepared protein

Plate 2. Prepared protein

Plate 1 is the original picture of the inflammatory protein as retrieved from protein databank (<http://www.rcsb.org>) with their co-crystallized ligands, which were used to validate the docking protocols for the binding sites while plate 2 is the prepared proteins as used for the docking, it was prepared by removing water of crystallization and unwanted protein chains in Discovery studio.

3. RESULTS AND DISCUSSION

3.1 Chemical Composition of the Seed Extract

The GC–MS analysis of Country Onion seed extract gave 18 peaks for the bioactive compounds, their percentage composition, retention time, structures and formulas were recorded in the Table 1.

The Table 1 showed the GC-MS result of the phytochemicals from ethanol seed extract of *A. lepidophyllus* and some commercial drugs.

The phytochemical components were presented in the order from the highest to the least percentage as follows: Eugenol (47.14) > 3-allyl-6-methoxyphenol (24.23) > caryophyllene (10.35) > Oleic Acid (7.91), > tricosane (2.34), > Cis- Vaccenic acid 2.03) > Phenol,2-methoxy-4-(2-propenyl)-,acetate (1.44), > Oleic acid (1.21), the other compounds present have percentage occurrence of less than 1.00. Eugenol, 3-allyl-6-methoxyphenol and caryophyllene have over 81% of the percentage composition of the plant.








Eugenol is the highest occurring compound in clove seed. Eugenol (EUG) is a versatile naturally occurring molecule as phenolic monoterpenoid and frequently found in essential

oils in a wide range of plant species. It has huge industrial applications particularly in cosmetics, dentistry, flavoring of foods, agriculture, and pharmaceuticals. Recently Eugenol has been a target due to its great potential in preventing several chronic conditions. The World Health Organization (WHO) has declared EUG as nonmutant and generally recognized as safe (GRAS) molecule. The available literature about pharmacological activities of EUG shows remarkable anti-inflammatory, antioxidant, antifungal, antibacterial, antipyretic activity, neurodegenerative disorders, analgesic, anti-diabetic, anticancer, cardiovascular Protection and Anti-parasite activity having significant effect on human health[13].

3-allyl-6-methoxyphenol is another major bioactive component present in the seed extract of country onions with percentage composition 24.23. It has wide range of bioactive activities such as anti-inflammatory, antiageing, antioxidant, antimicrobial and anti-cancer properties [14].

Caryophyllene is a natural bicyclic sesquiterpenoid abundantly found in essential oils from various spices, fruits and medicine as well as ornamental plants. It is approved by United States Food and Drug Administration and European agencies as food additive, taste enhancer and flavoring agent and as such termed phyto-cannabinoid. Various pharmacological activities such as cardio-protective, hepato-protective, gastro-protective, neuro-protective, nephron-protective, antioxidant, anti-inflammatory, antimicrobial and immune-modulator have been reported in experimental studies which also showed potent therapeutic promise in neuropathic pain, neurodegenerative and metabolic diseases [15].

Table 1. Result of GCMS Analysis of phytochemicals from Country Onion ethanol seed extract

S/N	RT	Phytochemicals	Percentage	Pubchem CID	Structure/Molecular formula
1	7.076	D-Limonene	0.28	440917	
2	16.69	Eugenol	47.14	3314	<p>$C_{10}H_{16}$</p> 
3	17.01	3-allyl-6-methoxyphenol	24.23	596375	<p>$C_{10}H_{12}O_2$</p> 
4	17.86	Bicyclo[7.2.0]undec-4-ene	0.47	129848326	<p>$C_{10}H_{12}O_2$</p> 
5	18.15	Caryophyllene	10.35	5281515	<p>$C_{11}H_{18}$</p> 
6	19.12	Humulene	0.58	5281520	<p>$C_{15}H_{24}$</p> 
7	20.83	2,4,5,7-tetrathiooctane	0.42	158825	<p>$C_{15}H_{24}$</p> 

S/N	RT	Phytochemicals	Percentage	Pubchem CID	Structure/Molecular formula
8	20.96	Phenol,2-methoxy-4-(2-propenyl)-,acetate	1.44	7136	$C_{15}H_{18}O_4$
9	22.44	Caryophyllene oxide	0.21	1742210	$C_{15}H_{14}O_3$
10	25.15	2,3,4-trimethoxyacetophenone	0.60	83810	$C_{15}H_{24}O_7$
11	29.51	Hexadecanoic acid, methyl ester	0.42	8181	$C_{17}H_{34}O_2$
13	29.99	Dibutyl phthalate	0.24	3026	$C_{17}H_{34}O_4$
14	31.13	11-octadecenoic acid, methyl ester	0.15	5364432	$C_{19}H_{36}O_2$
15	33.89	Tricosane	2.34	12534	$C_{23}H_{48}$
16	34.71	Oleic Acid	7.91	445639	$C_{18}H_{34}O_2$




S/N	RT	Phytochemicals	Percentage	Pubchem CID	Structure/Molecular formula
17	34.79	Cis- Vaccenic acid	2.03	5282761	 C ₁₈ H ₃₄ O ₂
18	36.24	9-Octadecenoic acid	0.81	637517	 C ₁₈ H ₃₄ O ₂
19	36.52	2-methyl-Z,Z-3,13-octadecadienol	0.20	5364412	 C ₁₉ H ₃₆ O

Table 2. Result of molecular docking of the country onion compounds, some commercial drugs with inflammatory protein (1pxx)

S/N	Phytochemicals	Percentage	Pubchem CID	Binding affinity(kcal/mol)
1	D-Limonene	0.28	440917	-6.5
2	Eugenol	47.14	3314	-6.5
3	3-allyl-6-methoxyphenol	24.23	596375	-6.1
4	Bicyclo[7.2.0]undec-4-ene	0.47	129848326	-7.4
5	Caryophyllene	10.35	5281515	-7.5
6	Humulene	0.58	5281520	-7.1
7	2,4,5,7-tetrathiooctane	0.42	158825	-3.5
8	Phenol,2-methoxy-4-(2-propenyl)-,acetate	1.44	7136	-6.5
9	Caryophyllene oxide	0.21	1742210	-7.2
10	2,3,4-trimethoxyacetophenone	0.60	83810	-6.2
11	Hexadecanoic acid, methyl ester	0.42	8181	-6.5
12	Oleic acid	7.91	445639	-6.7
13	Dibutyl phthalate	0.24	3026	-7.2
14	11-octadecenoic acid, methyl ester	0.15	5364432	-6.9
15	Cis- Vaccenic acid	2.03	5282761	-6.8
16	9-Octadecenoic acid	0.81	637517	-7.0
17	2-methyl-Z,Z-3,13-octadecadienol	0.20	5364412	-6.9
18	Ibuprofen		3672	-7.3
19	Naproxen		156391	-8.1
20	Indomethacin		3715	-7.7
21	Diclofenac		3033	-8.5

Humulene, also known as α -caryophyllene, is a ring-opened isomer of β -caryophyllene found in cannabis, ginseng and sage, Humulene possesses both topical and systemic anti-inflammatory properties, is an effective analgesic when taken topically, orally, or by aerosol, Interestingly, humulene was shown to increase secretion of IL-8, a chemokine with various functions, including promoting angiogenesis, helpful in wound healing [16]

Vaccenic acid is a naturally occurring trans fatty acid and an omega-7 fatty acid. It is the predominant kind of trans-fatty acid found in

human milk, in the fat of ruminants, and in dairy products such as milk, butter, and yogurt. cis-Vaccenic acid is used as a therapeutic agent against cardiovascular diseases, it has anticarcinogenic properties [17], and also helps in reduction of total cholesterol, LDL cholesterol and triglyceride level [18,19,20]

The 18 compounds from the seed extracts and some commercial drugs were docked on the protein to know their binding affinities and compared with that of some known anti-inflammatory drugs to determine the one with better binding affinity against the binding site of the disease protein.

The docking result showed that the drug Diclofenac has the best binding affinity of -8.5 kcal/mol followed by another drug Naproxen and indomethacin -8.1 and 7.7 kcal/mol respectively. the seed bioactive compound Caryophyllene, Bicyclo [7.2.0] undec-4-ene, Caryophyllene oxide, Dibutyl phthalate, and Humulene have binding affinities of -7.5, -7.4, -7.2 and -7.1 kcal/mol respectively higher and very close to that of a commercial drug Ibuprofen (-7.3 kcal/mol). Other compounds from the seed with good binding affinities between -6.1 to -6.9 kcal/mol, they include 9-octadecenoic acid (-7.0 kcal/mol), 11-octadecenoic acid, methyl ester (-6.9 kcal/mol), Phenol, 2-methoxy-4-(2-propenyl)-, acetate (-6.1 kcal/mol) except 2,3,5,7-

tetrathioactane (-3.5 kcal/mol). The interaction of the compounds with better binding affinities and the commercial drugs were carried out on the enzyme to confirm whether the docking was on the active site and pocket. The results were discussed below.

3.2 Interactions

Figs. 1-7 were the interaction results of the extracts with better binding affinities, and commercial drugs to determine the types of bond they formed with the disease protein, these were done using pyRx software and the results were recorded in Table 3.




Fig. 1. Interaction of the protein with Bicyclo[7.2.0]undec-4-ene.




Fig. 2. Interaction of the protein with caryophyllene




Fig. 3. Interaction of the protein with Dibutyl phthalate




Fig. 4. Interaction of the protein with Diclofenac




Fig. 5. Interaction of the protein with Ibuprofen




Fig. 6. Interaction of the protein with indomethacin




Fig. 7. Interaction of the protein with Naproxen

The interaction results above showed that some of the identified compounds have significant potentials to block the active site of the enzyme. Out of the 18 identified compounds, 3 compounds with binding affinities close to the reference drugs and the reference drugs were used to study their interaction with the disease protein, the drug Diclofenac has the minimum value of the binding affinity -8.5 and it interacted with the COX enzyme through two conventional hydrogen bond TYR 2385 and SER 2530, Pi sulfur bond with MET 2522, Pi-Pi T Shaped bond with TRP 2382, it also has pi alkyl bond with ALA 2527 and a hydrophilic Vander waals bond with LEU 2531, PHE 2518, ARG 2120, TYR 2355, SER2353, LEU 2534, LEU 2352, PHE 2205,

TYR 2348, PHE 2381, GLY2526 and LEU 2384. Naproxen another commercial drug with affinity - 8.1kcal/mol displayed the strongest interaction with the diseased protein among all docked ligand in this study, Naproxen constructed a carbon hydrogen bond with ARG 2120 and a conventional hydrogen bond with GLY 2526. Pi sigma bond with VAL 2349, ALA 2527, Pi Alkyl with LEU 2531, PHE 2351, VAL 2523 and TUR 2385 and a hydrophobic vander waal bond with MET 2113, TYR 2355, VAL 2116, LEU 2359, SER 2359, SER 2353, SER 2530, TRP 2387, LEU 2352. Indomethacin another control drugs showed a strong carbon hydrogen bond and pi donor hydrogen bond with PHE 2518, GLY 2526 and SER 2353 amino acids, It also has pi sigma

bond with VAL 2523, pi alkyl and alkyl bond with HIS 2090, LEU 2352, LEU 2359, TYR 2355, VAL 2116, LEU 2531 and VAL 2349, and vander waal interaction with MET 2522, TRP 2387, LEU 2384, PHE 2381, SER 2530, ARG 2120, ARG 2513. The high binding affinity may be due to its strong binding with other residue at the active site. Dibutyl phythate although not the closest ligand in binding affinity to the contro drug showed a similar type of bond with Naproxen and best interaction among the phytocompounds docked having a conventional hydrogen bond with ARG 2120, a carbon hydrogen bond with ALA 2527 and SER 2353, it has Pi sigma bond

with Val 2349 and Pi Alkyl and Alkyl bonds with LEU 2359, LEU 2531, VAL 2116, LEU 2352 and Val 2523. It also has hydrophobic Vander waal bond with amino acids MET 2113, TRR 2355, PHE 2518, SER 2530. GLY 2526, PHE 2382, TYR 2385, TRP 2387 and TYR 2348. Caryophyllene and Bicyclo [7.2.0]undec-4-ene with binding affinities -7.5 and -7.4 kcal/mol have most of their interactions with hydrophobic vander waals and alkyl bonds, however did not construct strong bonds with the major active residue instead have a weak interaction. These compounds serve as leads in treating inflammatory diseases.

Table 3. Docking results showing the type of interaction and amino acid involved

Compound	Binding affinity	Type of interaction	Amino acids
Bicyclo [7.2.0]undec-4-ene	-7.4	Vander waals	ALA 2527, SER 2530, TYR 2348, GLY2526 GLY2526, PHE 2381, LEU 2384, MET 2522, PHE 2518
		Alkyl and pi-Alkyl	LEU 2354, TYR 2385, VAL 2349, TRP 2387 VAL 2523,
Caryophyllene	-7.0	Vander waal	ARG 2120, SER 2353, TYR 2355, PHE 2518, LEU2352, GLU 2526, SER 2530
		Alkyl	LEU 2531, LEU 2389, VAL 2349, VAL2523, ALA2527
Dibutyl phthate	-7.2	Vander waal	TYR 2355, MET 2113, PHE 2518, SER 2530 GLY 2521, PHE 2385, TYR 2385, TRP 2387 TYR 2348
		Conventional hydrogen bond	ARG 2120
		Carbon hydrogen bond	ALA 2527, SER 2353
		Pi sigma bond	VAL 2349
		Alkyl and Pi- alkyl	VAL 2523, LEU 2352 LEU 2331, VAL 2116 LEU 2359
Diclofenac	-8.5	Vander waal	LEU 2531, PHE 2518 ARG 2120, TRY 2355 SER 2353, LEU 2534 LEU 2352, PHE 2205 TYR 2348, PHE 2381 GLY 2526, LEU 2384 TYR 2385, SER 2530 MET 2522
		Conventional hydrogen bond	MET 2522
		Pi-Sulfur	TRP 2387
		Pi-Pi T shaped	ALA 2527, VAL 2523 VAL 2349
Naproxen	-8.1	Pi alkyl	
		Vander waal	MET 2113, TYR 2355 VAL 2116, SER 2353, TRP 2387, SER 2530 LEU 2359
		Conventional hydrogen bond	ARG 2120
		Carbon hydrogen bond	
		Pi-sigma bond	GLY 2526

Compound	Binding affinity	Type of interaction	Amino acids
Indomethacin	-7.7	Pi Alkyl	VAL 2349, ALA 2527 PHE 2381, VAL 2523 TYR 2385, LEU 2531
		Vander waal	MET 2522, TRP 2387 LEU 2384, PHE 2381 SER 2530, ARG 2120, ARG2513, PHE 2518, GLY 2526, SER 2353
		Pi- sigma bond	VAL 2523
Ibuprofen	-7.3	Alkyl	LEU 2531, VAL 2116, TYR 2355, LEU 2359
		Vander waal	ALA 2527., HIS 2090 LEU 2352, VAL 2349 ARG 2120, SER 2530, PHE 2381, GLY 2526, TYR 2385 LEU 2352, VAL 2523 TYR 2348 , SER 2353, LEU 2531, MET 2113
		Pi – sigma	VAL 2349, ALA 2527 VAL 2116, LEU 2359. TYR 2355
		Alkyl and Pi alkyl	

4. CONCLUSION

The result obtained from the GC-MS showed that *Afrostryrax lepidophyllus* seed contain 18 bio-active compounds with known medicinal, biological and therapeutic properties with their percentage composition as follows: Eugenol (47.14%), 3-Ally-6-methoxyphenol (24.23%), Carophyllene (10.35%) and Oleic acid (7.91%) were the most abundant compounds in the extract. Eugenol and Carophyllene both amounting to 57.49% of the total percentage of compound found in the extract have been reported to show anti-inflammatory activity thereby positively validating the claims made by the Cameroonians that *Afrostryrax lepidophyllus* can cure inflammatory diseases [21]. Inflammation has been an underlying factor in most health condition like diabetes, arthritis, heart disease and cancer. The dried seed of *A. lepidophyllus* is reported to have been used medically to reduce inflammation. This can be attributed to the presence of caryophyllene, eugenol humulene, Dibutyl phythate etc. *A. lepidophyllus* seed contains antimicrobial, antioxidant and larvidival compounds such as M. eugenol, tricosane and oleic acid (anticancer). Also, compounds like Cis-vaccenic acid and D-limonene present in the extract were reported to reduce cholesterol and prevent heart diseases [8]. Humulene was reported to have analgesic property (pain killer) [22], It also possesses carminative properties that help to reduce flatulence and stomach gas [23]. The use of *A. lepidophyllus* seed to treat high blood sugar level in people with diabetes can be attributed to the

presence of cis-vaccenic acid. The research revealed that the efficacy of *A. lepidophyllus* seed medicinally is due to its phytochemicals [24] and that the seed potentially aid inflammation reduction and associated diseases as recorded by previous researchers [8]. Further studies needs to be carried out using molecular dynamic simulation to determine the stability of these extracts (ligands) towards the protein moiety over time.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dini I. Spices and herbs as therapeutic foods. In: Food quality: Balancing health and disease. Cambridge, Massachusetts, United States: Academic Press. 2018;433-469.
2. Jiang TA. Health benefits of culinary herbs and spices. Journal of AOAC International. 2019;102(2):395-411. DOI:10.5740/jaoacint.18-0418.PMID: 30651162

3. Hossain MS, Uddin MS, Kabir MT, Akhter S, Goswami S, Mamun AA, Herrera-Calderon O, Asaduzzaman M, Abdel-Daim MM. *In vivo* Screening for analgesic and anti-inflammatory activities of *Syngonium podophyllum* L.: A remarkable herbal medicine. Ann. Res. Rev. Biol. 2017, Sep. 2;16(3):1-12. [cited 2024 May 31] Available:<https://journalarrb.com/index.php/ARRB/article/view/1101>
4. Ify OA, Raphael AG, Tochukwu OC, Amarachi OUS, Ikechukwu NA, Madukaihe MJ, Thomas Y, Innocent OC. The antimicrobial, anti inflammatory and analgesic activities of the rhizome extract of *Curcuma longa* L. (Turmeric). J. Adv. Biol. Biotechnol. 2021, Jul. 12;24(6):1-16. [cited 2024 May 31] Available:<https://journaljabb.com/index.php/JABB/article/view/502>
5. Ngenge TA, Jabeen A, Maurice TF, Baig TA, Shaheen F. Organic and mineral composition of seeds of *Afrotyrax lepidophyllus* Mildbr. and evaluation of ROS inhibition and cytotoxicity of isolated compounds. Chemistry Africa. 2019, Dec;2:615-24.
6. Embuscado ME. Spices and herbs: Natural sources of antioxidants—A mini review. Journal of Functional Foods. 2015;8:11-819.
7. Jun LV. Consumption of spicy foods and total and cause specific mortality: Population based cohort study. BMJ; 2015. DOI: 10.1136/bmj.h3942 Available:<https://www.sciencedaily.com › releases › 2015/08>
8. Smoky Hazel, Surprising Health Benefits of Country Onions You Never Knew Available:<https://smokyhazelspice.com › blogs › surprising-health...>
9. Rhiannon CT. Country Onion” (*Afrotyrax lepidophyllus*): Spice and medicine of the cameroon rainforest, Conference: 94th ESA Annual Convention; 2009.
10. Nortjie E, Basitere M, Moyo D, Nyamukamba P. Extraction methods, quantitative and qualitative phytochemical screening of medicinal plants for antimicrobial textiles: A Review. Plants (Basel). 2022;11(15): 2011. DOI:10.3390/plants11152011PMCID: PMC9370299
11. Duru CE, Duru IA, Adegboyega AE. *In silico* identification of compounds from *Nigella sativa* seed oil as potential inhibitors of SARS-CoV-2 targets Bull Natl Res Cent. 2021;45(57):1-13. Available:<https://doi.org/10.1186/s42269-021-00517-x>
12. Ugariogu SN, Duru IA. Anticolorectal Cancer Properties of Some Flavonoids and Terpenoids from African Propolis via Molecular Docking South Asian Research Journal of Natural Products. 2022;5(3): 209-221.
13. Muhammad FN, Mahnoor K, Muhammad R, Jinyin C, Yali Y, Chunpeng CW. Pharmacological properties and health benefits of eugenol: A comprehensive review. Hindawi Oxidative Medicine and Cellular Longevity; 2021.
14. Orlo E, Russo C, Nugnes R, Lavorgna M, Isidori M. Natural methoxyphenol compounds: Antimicrobial activity against food borne pathogens and food spoilage bacteria, and role in antioxidant processes. 2021;10(8):1807. DOI:10.3390/foods10081807.
15. Sharma C, Al Kaabi JM, Nurulain SM, Goyal SN, Kamal MA, Ojha S. Polypharmacological properties and therapeutic potential of β -caryophyllene: a dietary phytocannabinoid of pharmaceutical promise. Curr Pharm Des. 2016;22(21):3237-3264. DOI: 10.2174/1381612822666160311115226
16. Gupta CR. Humulene, Nutraceuticals, Efficacy, safety and toxicity, Academic press. 2016;983-1027
17. Lock AL, Corl BA, Barbano DM, Bauman DE. The anticarcinogenic effect of trans-11 18:1 is dependent on its conversion to cis-9, trans-11 CLA by delta9-desaturase in rats. J Nutr. 2004;134(10):2698–704. DOI:10.1093/jn/134.10.2698. PMID 15465769.
18. Vaccenic Acid. Jahrels G, Dawczyask C. Milk and Dairy food; 2020. Available:<http://www.science direct.com>
19. Linlin C, Huidan H, Cui H, Jing J, Zuo, Deng, J, Li Y, Wang X, Zhao L. Inflammatory responses and inflammation-associated diseases in organs. Oncotarget. 2018;9(6):7204–7218. DOI: 10.18632/oncotarget.23208
20. Ricciotti E, FitzGerald GA. Prostaglandins and Inflammation, Arterioscler Thromb Vasc Biol. 2012;31(5):986–1000.

21. Takechi M, Matsunami S, Nishizawa J, Uno C, Tanaka Y. Haemolytic and antifungal activities of saponins or anti-ATPase and antiviral activities of cardiac glycosides *planta Medica*. 1999;(65):585-586.
22. Russo EB, Jahan Marcu. Cannabis pharmacology: The usual suspects and a few promising leads advances in pharmacology. 2017;80(2):67-134.
23. Dahee Y, Su-Jung H, Ye-Seul S, Hyo-Jong L. Humulene Inhibits Acute Gastric Mucosal Injury by Enhancing Mucosal Integrity Antioxidants (Basel). 2021 May; 10(5):761.
DOI:10.3390/antiox10050761PMCID: PMC8150829
24. Yilma HG, Fekade BT, Archana B, Nishant R, Mesfin GT, A Nasser, S Kundan, KC, Rakesh KB. Anti-inflammatory activity of phytochemicals from medicinal plants and their nanoparticles: A review. *Current Research in Biotechnology*. 2023;6: 100152.
Available:<https://doi.org/10.1016/j.crbiot.2023.100152>Get rights and content

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/118460>