



Effect of NPK, Zinc and Iron on Soil Health and Yield Parameters of Green Gram (*Vigna radiata* L.) var SML-668

Prerna Timsina ^{a++}, Arun Alfred David ^{a#},
Narendra Swaroop ^{a#}, Tarence Thomas ^{a#*},
Ram Bharose ^{a†} and Jagdish Saran ^{a++}

^a Department of Soil Science and Agricultural Chemistry, [Naini Agricultural Institute], Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj- 211 007, (Uttar Pradesh), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71036>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118530>

Original Research Article

Received: 16/04/2024
Accepted: 18/06/2024
Published: 22/06/2024

ABSTRACT

A study on the effects of NPK, zinc, and iron on soil health and yield parameters of green gram was carried out in the crop research farm, department of Soil Science and Agricultural Chemistry at SHUATS, Prayagraj, from May to July of 2023. Nine treatments were used in a randomised block design (RBD) experiment, each of which was replicated three times. During the sowing process,

⁺⁺ Research Scholar;

[#] Associate Professor;

[†] Assistant Professor;

*Corresponding author: E-mail: prernatimsina929@gmail.com;

Cite as: Timsina, Prerna, Arun Alfred David, Narendra Swaroop, Tarence Thomas, Ram Bharose, and Jagdish Saran. 2024. "Effect of NPK, Zinc and Iron on Soil Health and Yield Parameters of Green Gram (*Vigna Radiata* L.) Var SML-668". *Journal of Advances in Biology & Biotechnology* 27 (7):769-80. <https://doi.org/10.9734/jabb/2024/v27i71036>.

NPK fertiliser, zinc, and iron were added to the soil. The recommended doses of zinc and iron are 15 kg ha⁻¹ and 20 kg ha⁻¹, respectively, and of NPK are 25:50:25. The plant height, No. of pods plant⁻¹, No. of seeds pod⁻¹, grain yield, water holding capacity, % pore space, % organic carbon, soil EC, available nitrogen, available potassium, and available iron were found to be significant and was recorded maximum in T₉- [[NPK @100% + Zn @100% + Fe @100%]. Bulk density, particle density and soil pH were recorded maximum in T₁- [NPK @ 0 % + Zn @ 0 %+ Fe @ 0%]. Available phosphorous was recorded maximum in T₇- [NPK @100% + Zn @0% + Fe @100%] whereas available zinc was recorded maximum in T₃- [NPK @ 0% + Zn @ 100% + Fe @100%]. The present study reported that the Zinc and Iron on Soil Health and Yield Parameters of Green gram (*Vigna radiata* L.) var. SML-668 fertilizes doses of NPK + Zinc + Iron gave positive test values arching desired yield targets of greengram situations for Prayagraj region.

Keywords: Soil health; yield parameters; zinc; iron; green gram.

1. INTRODUCTION

Green gram is grown best in the alluvial tract in the north, as well as black and red soils of southern and peninsular India. It can also be grown on light stony soil to clayey soils. Green gram cultivation is ideal in well drained loam or sandy loam soils. In India during 2023-24, about 15.93 lakh ha (39.38 lakh acres) area was covered under green gram as against 15.57 lakh ha (38.47 lakh acres) during the same period in 2022-23. According to Government 3rd advance estimates, green gram production in 2022-23 was 3.74 million tonnes. The highest production was reported in Rajasthan (39.06%), followed by Madhya Pradesh (11.57%), Maharashtra (8.57%), Bihar (5.54%), Karnataka (5.05%), Tamil Nadu (5.02%), Gujarat (4.63%), and Andhra Pradesh (4.31%) [1]. According to directorate of economics and statistics, DAC and FW the total production of green gram in Uttar Pradesh in the year 2020-21 was 61 tonnes ha⁻¹. India is the world's top producer of green gram, it is farmed on roughly 4.5 million hectares, producing 2.5 million tonnes at a productivity of 548 kg per hectare, making up 10% of the world's production of pulses. Green gram is grown in almost all the states in India. Green gram is the most important crop of the south-east Asia and the most importantly of Indian sub-continent [2]. Being a leguminous crop, green gram has the capacity to fix the atmospheric nitrogen. Due to its less water requirement and deep rooting system green gram is favourable in semi-arid tropics. When growth conditions are suitable, greengram, often known as mung beans, go through an epigeal germination process that takes 4-5 days [3-5]. The plant has numerous lateral roots and a well-developed root system with root nodules. Green gram is a short duration crop. Being a short duration crop it fits well in many intensive crop rotations. Green

gram can be grown thrice a year, in the Kharif season during July to October, in the Rabi season during November to March and Summer season during April to June [6]. Green gramme is known as the "green pearl" because of its high nutrient content [7]. Green gram is consumed in different forms such as vegetables, sprouts, dhal, processed grain, fried bean, bean paste or incorporated into noodles, bread, cakes, cold jellies, and desserts [8]. India is the largest producer of green gram and account for 54% of the world production and covers 65% of the world acreage. Besides being the largest pulses producer, India is also one of the largest consumers and importers in the world, because of the rise in population and consumer awareness, India imports pulses to meet the high domestic demand (Pulses revolution from food to nutritional security, success report, 2017-18). Green gram is also known as poor man's meat [9]. By improving the physical, biological, and chemical characteristics of the soil as well as its fertility status through biological nitrogen fixation in a symbiotic relationship with rhizobium from the atmosphere and by mushrooming the number of soil microorganisms, pulses enhance soil health [10]. Mung bean contains about 51.6% carbohydrate, 26 to 27% protein, 4 to 5% minerals and 3 to 4% vitamins [11].

Among all the macronutrients that are typically applied as commercial fertilisers, nitrogen is the first ingredient. Given that it appears to have the most immediate and noticeable effect, nitrogen is one of the most vital nutrients for plants. Nitrogen gives plants their dark colour and promotes their above-ground vegetative growth. In plants, nitrogen plays a unique role in the synthesis of protein. A lack of nitrogen might cause plants to develop more slowly.

The second fertiliser ingredient is phosphorus, which is necessary for the nourishment of plants as well as for every living cell. It participates in every kind of plant metabolism. It is a structural element of the cell membrane system and a necessary component of most enzymes. Root growth is stimulated by phosphorous. It stimulates rhizobial activity and the production of crop nodules.

Potassium is the third fertilizer element. Potassium acts as a root booster, stalk strengthner, food former, sugar and starch supporter. It is also essential for the photosynthesis. It increases the boldness of the grain. Potassium is also refered to as a value element because of its contributions to the quality, taste, colour etc [12].

Zinc is a trace element, it is very significant as it performs numerous functions such as proper growth of the plant, synthesis of chlorophyll, biosynthesis of plant growth hormone and RNA synthesis. Green gram when grown under zinc deficient soils suffers from yield loss [13]. Zinc insufficiency is the most often reported deficit. Zinc shortage has been reported in many states of India (Likhittakutum et al. 2023).

Whereas iron plays a very important role in nitrogen fixation. It regulates respiration, photosynthesis, Sulphur absorption and nitrogen-fixing. It is also involved in the synthesis of protein. Therefore, present investigation was undertaken to study the effect of NPK, Zinc and Iron on soil health parameters and yield.

2. MATERIALS AND METHODS

The soil samples were collected randomly from the experimental site prior to tillage operation as well as after the harvest from a depth of 0-15 and 15-30 cm. The topography of experimental field was uniform and levelled. The experimental soil was sandy loam in texture. Seed was sown by dibbling method with the spacing of 30 cm between rows and 10 cm within plants. The experimental field was well prepared by ploughing with tractor. The field was leveled and stubbles and weed were picked up from the field manually. As per experimental recommendations, the fertilizers Nitrogen, Potassium, Phosphorous, Zinc and Iron were weighed and applied in the field by mixing thoroughly with soil. Thinning was done to maintain proper spacing of 10 cm between plants. In order to keep field free from weeds, and to ensure soil moisture two hand weeding was done at 15 and 35 DAS to maintain the optimum population of the plant.

Physical parameters like bulk density, particle density, pore space and water holding capacity was analysed through 100 ml graduated measuring cylinder method and the process given by Muthuval et al. [14].

Soil chemical parameters:

- a. Soil pH – [15]
- b. Soil EC (dS m^{-1}) – [16]
- c. Organic carbon (%) – [17].
- d. Available Nitrogen (kg ha^{-1}) – [18].
- e. Available Phosphorus (kg ha^{-1}) – [19].
- f. Available Potassium (kg ha^{-1}) – [20].
- g. Available Zinc and Iron (Mg kg^{-1}) – [21].

Table 1. Treatment combinations of Greengram

Treatment	Treatment Combination
T ₁	[NPK @ 0% + Zn @ 0% + Fe @ 0%]
T ₂	[NPK @ 0% + Zn @ 50% + Fe @ 100%]
T ₃	[NPK @ 0% + Zn @100% + Fe @ 100%]
T ₄	[NPK @ 50% + Zn @ 0% + Fe @ 100%]
T ₅	[NPK @ 50% + Zn @ 50% + Fe @100%]
T ₆	[NPK @ 50% + Zn @ 100% + Fe @ 100%]
T ₇	[NPK @ 100% + Zn @ 0 % + Fe @ 100 %]
T ₈	[NPK @100% + Zn @ 50% + Fe @ 100 %]
T ₉	[NPK @100 % + Zn @ 100% + Fe @ 100 %]

3. RESULTS AND DISCUSSION

3.1 Physical Properties of Soil

As depicted in Table 2. the soil bulk density, particle density, water holding capacity and pore space was found non-significant. Bulk density and particle density increases with the increase in depth. Similar findings were recorded by Chintha et al. [22] and Hussain et al. [23]. Whereas water holding capacity and pore space was found significant. Water holding capacity increases with the increase in depth, the maximum data recorded was 45.31% and 45.71 % in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by 45.25% and 45.64 % in T₅ [NPK @50% + Zn @50% + Fe @100%] and minimum was 43.13 % and 43.21% in T₁. [NPK @ 0% + Zn 0% + Fe @ 0 %]. Similar findings were recorded by Prakash et al. [24], Banjara et al. [25]. Soil pore space decreases with the decrease in depth. The maximum value recorded for pore space was 48.76% and 48.53 % at 0-15 cm and 15-30 cm respectively in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by 48.60% and 48.48 % in T₅ [NPK @50% + Zn @50% + Fe @100%] and the minimum value recorded was 46.21% and 46.15% in T₁ [NPK @0% + Zn @0% + Fe @0%]. Similar findings were recorded by Tarun et al. [26].

3.2 Chemical Properties of Soil

As depicted in Table 3. the value of soil pH, soil EC and organic carbon, Soil pH and soil EC was found non-significant whereas organic carbon was found to be significant. The maximum pH of soil 7.25 and 7.37 was found in treatment T₁ [NPK @ 0 % + Zn @ 0 % + Fe @0%] and the minimum pH of soil 6.85 and 6.92 was found in treatment T₉ [NPK @100% + Zn @100% + Fe @100%]. Similar findings were recorded by Lokendra et al. [27]. The maximum EC value 0.36 dSm⁻¹ and 0.40 dSm⁻¹ was recorded in T₉ [NPK @100% + Zn @100% + Fe @100%]. The minimum EC value 0.21 dSm⁻¹ and 0.23 dSm⁻¹ was recorded in T₁ [NPK @0% + Zn @0% + Fe @0%]. The maximum value of EC in T₉ might be due to the application of 100% inorganic fertilizers which results in an increase in salt content in soil, as soil EC is directly proportional to the nutrient concentration level, and inversely proportional to the depth. Similar findings were recorded by Likhithakuttum et al. (2022) and Lokendra et al. [27]. The maximum organic carbon of soil 0.55% and 0.57 % was found in treatment T₉ [NPK @100% + Zn @100% + Fe

@100%] and minimum organic carbon of soil 0.30% and 0.28 % was found in treatment T₁ [NPK @0% + Zn @0% + Fe @0%] respectively. The increased organic carbon (%) might be due to the fertilization which indirectly increases the soil organic carbon. Inorganic fertilizers improve the soil organic matter content in the soil by increasing the plant biomass which remains in the field and undergoes decomposition thus increasing the soil organic matter. Similar findings were recorded by Sahu et al. [28], Deshlahare et al. [29]. As depicted in the Table 4 and Fig 1. the value of NPK was found to be significant. The maximum available nitrogen 304.19 Kg ha⁻¹ and 306.20 kg ha⁻¹ was recorded maximum in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by 303.20 Kg ha⁻¹ and 304.99 kg ha⁻¹ was found in T₇ [NPK @100% + Zn @0% + Fe @100%] and the minimum value recorded was 275.85 Kg ha⁻¹ and 276.98 kg ha⁻¹ was found in T₁ [NPK @0% + Zn @0% + Fe @0%]. The application of NPK together with Zinc and Iron resulted in significantly increase of nitrogen in soil, it might be due to increased microbial activity leading to the mineralization of nutrients. The increase in the nitrogen content may be due to the synergistic effect of Zinc on Nitrogen in soil. Similar findings were recorded by Hemraj et al. [30], Mohammad et al. [31]. The maximum available phosphorous 31.50 kg ha⁻¹ and 32.51 kg ha⁻¹ was recorded in T₇ [NPK @100% + Zn @0% + Fe @100%] followed by 30.87 and 31.64 kg ha⁻¹ in T₄ [NPK @50% + Zn @0% + Fe @100%] and the minimum value recorded was 26.48 Kg ha⁻¹ and 28.05 kg ha⁻¹ in T₁ [NPK @0% + Zn @0% + Fe @0%]. Phosphorous content increases with the increase in level of NPK whereas it decreased with an increase in level of Zinc due to its antagonist effect with zinc. Similar findings were recorded by Koushik et al. [32]. The maximum value of available potassium recorded was 210.07 kg ha⁻¹ and 213.27 kg ha⁻¹ in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by 207.99 kg ha⁻¹ and 211.4 kg ha⁻¹ in T₇ [NPK @100% + Zn @0% + Fe @100%] and the minimum available potassium recorded was 180.85 and 182.84 kg ha⁻¹ in T₁ [NPK @0% + Zn @0% + Fe @0%]. Similar findings were recorded by Mohammad et al. [31], Chethan KV et al. [33], Rohit et al. [34]. As depicted in Table 5 and Fig 2. the value of zinc and iron was found significant. The value of available zinc increases with the increase in depth. The maximum value of zinc recorded was 0.94 kg ha⁻¹ and 0.97 kg ha⁻¹ in T₃ [NPK @0% + Zn @100% + Fe @100%] followed by 0.91 kg ha⁻¹ and 0.89 kg ha⁻¹ in T₉ [NPK @100% + Zn @100% + Fe @100%] and

the minimum value recorded was 0.71 kg ha⁻¹ and 0.75 kg ha⁻¹ in T₁ [NPK @0% + Zn @0% + Fe @0%]. The minimum value of Zinc might be due to the reaction of zinc with phosphorous which results in the formation of insoluble compounds of Zn. Similar findings were recorded by Choudhary et al. [35], Rajendra et al. [36]. The value of available iron increase with the increase in depth. The maximum value of iron recorded was 6.59 kg ha⁻¹ and 6.84 kg ha⁻¹ in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by 6.47 kg ha⁻¹ and 6.78 kg ha⁻¹ in T₅ [NPK @50% + Zn @50% + Fe @100%] and the minimum value recorded was 5.10 kg ha⁻¹ and 5.07 kg ha⁻¹ in T₁ [NPK @0% + Zn @0% + Fe @0%]. Similar findings were recorded by Fulpagare et al. [37], Sunil et al. [38].

3.3 Growth Parameters

As depicted in Table 6. the value of plant height, No. of pod plant⁻¹, No. of seeds pod⁻¹ and grain yield are found to be significant. At 15,30,45 and 60 DAS the plant height was recorded maximum in T₉ [NPK @100% + Zn @ 100% + Fe@100%] Followed by T₅ [NPK@ 50% + Zn@ 50% + Fe@100%] and the minimum plant height was recorded in T₁ [NPK @ 0% + Zn@0% + Fe @0%]. The increase in plant height might be due to the role of zinc and iron in various physiological activities such as, enzyme activation, chlorophyll synthesis, photosynthesis, cell elongation and differentiation which resulted in the vigorous growth of plant. Similar findings were also reported by Tribhuwana et al. [39]. No of pods plant⁻¹ was recorded maximum in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by T₅[NPK @50% + Zn @50% + Fe @100%] and the minimum No. of pods plant⁻¹ was recorded in T₁[NPK @0% + Zn @0% + Fe

@0%]. An increase in the No. of pods plant⁻¹ might be due to an increase availability of Iron which helps in sufficient absorption of nutrients. Zinc also plays a very significant role in affecting the growth parameters as it has a direct influence on the synthesis of tryptophan which is a precursor for the production of growth hormones known as auxin. Similar findings were recorded by Maddila et al. [40] and Vinodkumar et al. [41]. The No of seeds pod⁻¹ was recorded maximum in T₉ [NPK @100% + Zn @100% + Fe @100%] followed by T₅ [NPK @50% + Zn @50% + Fe @100%] and the minimum No. of seeds pod⁻¹ was recorded in T₁ [NPK @0% + Zn @0% + Fe @0%]. The increase in No of seed pod⁻¹ might be due to the important role of iron and zinc in affecting the growth characters and also the yield attributes. Other reasons may be due to the involvement of Zinc in IAA synthesis, IAA is a hormone which plays a major role in preventing the pod abscission which determines the seed yield. Zinc also plays a major role in seed setting. Similar findings were recorded by Boradkar et al. [42]. The grain yield was recorded maximum in T₉ [NPK@100% + Zn @100% + Fe @100%] followed by T₅[NPK @50% + Zn @50% + Fe @100%] and the minimum grain yield was recorded in T₁ [NPK @0% + Zn @0% + Fe @0%] [43-46]. Increase in green gram grain yield might be due to enhancement of pod formation and increase in number of seeds pod⁻¹ [47-50]. Highest grain yield obtained maybe due to the availability of 100% Zinc and Iron at all the growth stages, which resulted in enhanced metabolic process of plant which resulted in better yield attributes and application of Zinc and Iron directly in the soil has a synergistic effect on yield component. Similar finding was recorded by Jyoti et al. [51], Partha et al. [13], Gaffar et al. [52].




Fig. 1. Effect of NPK, Zinc and Iron on available NPK of post- harvest soil

Table 2. Effect of NPK Zinc and Iron on bulk density (Mg m^{-3}) particle density (Mg m^{-3}) water holding capacity (%) and pore space (%) of post-harvest soil

Treatment	Treatment Combination	Bulk density (Mg m^{-3})		Particle density (Mg m^{-3})		Water holding capacity (%)		Pore space (%)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	NPK @0% + Zn @0% + Fe @0%	1.304	1.305	2.55	2.58	43.13	43.21	46.21	46.15
T ₂	NPK @0% + Zn @50% + Fe @ 100%	1.261	1.265	2.45	2.49	43.37	43.39	47.41	47.36
T ₃	NPK @0% + Zn @100% + Fe @100%	1.291	1.292	2.38	2.45	43.33	43.44	47.46	47.40
T ₄	NPK @50% + Zn @0% + Fe @100%	1.283	1.284	2.33	2.37	44.40	44.50	47.57	47.50
T ₅	NPK @50% + Zn @50% + Fe @100%	1.274	1.276	2.15	2.19	45.25	45.64	48.60	48.48
T ₆	NPK @50% + Zn @100% + Fe @100%	1.291	1.293	2.17	2.21	45.18	45.22	48.43	48.29
T ₇	NPK @100% + Zn @0% + Fe @100%	1.283	1.285	2.23	2.26	45.14	45.16	48.47	48.35
T ₈	NPK @100% + Zn @50% + Fe@100%	1.287	1.290	2.28	2.32	45.07	45.09	48.51	48.42
T ₉	NPK @100% + Zn @100% + Fe @100%	1.268	1.274	2.09	2.12	45.31	45.71	48.76	48.53
	F-test	NS	NS	NS	NS	S	S	S	S
	C.D. at 5%	0.032	0.029	0.341	0.301	0.135	0.068	0.151	0.059
	S.Ed. (+)	0.015	0.013	0.161	0.142	0.064	0.032	0.071	0.028

Table 3. Effect of NPK, Zinc and Iron on soil pH, soil EC (dS m⁻¹) and organic carbon (%) of post-harvest soil

Treatment	Treatment Combination	Soil pH		Soil EC (dS m ⁻¹)		organic carbon (%)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	NPK @0% + Zn @0% + Fe @0%	7.25	7.37	0.21	0.23	0.30	0.28
T ₂	NPK @0% + Zn @50% + Fe @ 100%	7.23	7.32	0.27	0.29	0.32	0.31
T ₃	NPK @0% + Zn @100% + Fe @100%	7.18	7.25	0.33	0.36	0.37	0.34
T ₄	NPK @50% + Zn @0% + Fe @100%	7.16	7.21	0.32	0.35	0.40	0.39
T ₅	NPK @50% + Zn @50% + Fe @100%	6.94	6.93	0.34	0.36	0.54	0.55
T ₆	NPK @50% + Zn @100% + Fe @100%	7.09	7.14	0.30	0.33	0.51	0.52
T ₇	NPK @100% + Zn @0% + Fe @100%	7.11	7.19	0.28	0.31	0.46	0.47
T ₈	NPK @100% + Zn @50% + Fe@100%	7.07	7.14	0.24	0.25	0.42	0.44
T ₉	NPK @100% + Zn @100% + Fe @100%	6.85	6.92	0.36	0.40	0.55	0.57
	F-test	NS	NS	NS	NS	S	S
	C.D. at 5%	0.294	0.380	0.112	0.131	0.02	0.018
	S. Ed. (+)	0.138	0.179	0.052	0.061	0.01	0.008

Table 4. Effect of NPK, Zinc and Iron on available nitrogen (kg ha⁻¹), phosphorous (kg ha⁻¹) and potassium (kg ha⁻¹) of post-harvest soil

Treatment No.	Treatment Combination	Available Nitrogen (kg ha ⁻¹)		Available phosphorous (kg ha ⁻¹)		Available potassium (kg ha ⁻¹)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	NPK @0% + Zn @0% + Fe @0%	275.85	276.98	26.48	28.05	180.85	182.84
T ₂	NPK @0% + Zn @50% + Fe @ 100%	281.14	283.98	27.34	27.64	186.20	186.79
T ₃	NPK @0% + Zn @100% + Fe @100%	294.00	287.31	27.55	27.77	191.17	192.40
T ₄	NPK @50% + Zn @0% + Fe @100%	290.87	293.40	30.87	31.64	192.57	194.00
T ₅	NPK @50% + Zn @50% + Fe @100%	295.64	296.86	28.54	29.43	198.70	201.55
T ₆	NPK @50% + Zn @100% + Fe @100%	285.12	294.49	29.07	29.24	202.18	203.29
T ₇	NPK @100% + Zn @0% + Fe @100%	303.20	304.99	31.50	32.51	207.99	211.49
T ₈	NPK @100% + Zn @50% + Fe@100%	298.07	303.58	29.75	30.27	205.99	207.18
T ₉	NPK @100% + Zn @100% + Fe @100%	304.19	306.20	28.23	28.73	210.07	213.37
	F-test	S	S	S	S	S	S
	C.D. at 5%	1.511	1.976	0.364	0.823	1.945	1.243
	S. Ed. (+)	0.713	0.932	0.171	0.388	0.917	0.586

Table 5. Effect of NPK, Zinc and Iron on available Zinc (Mg kg⁻¹) and iron (Mg kg⁻¹) of post-harvest soil

Treatment No.	Treatment Combination	Available Zinc (Mg kg ⁻¹)		Available Iron (Mg kg ⁻¹)	
		0-15 cm	15-30 cm	0-15 cm	15-30 cm
T1	NPK @0% + Zn @0% + Fe @0%	0.71	0.75	5.10	5.07
T2	NPK @0% + Zn @50% + Fe @ 100%	0.76	0.81	5.63	5.72
T3	NPK @0% + Zn @100% + Fe @100%	0.94	0.97	5.75	5.83
T4	NPK @50% + Zn @0% + Fe @100%	0.83	0.85	5.84	5.88
T5	NPK @50% + Zn @50% + Fe @100%	0.86	0.92	6.47	6.78
T6	NPK @50% + Zn @100% + Fe @100%	0.89	0.94	6.09	6.14
T7	NPK @100% + Zn @0% + Fe @100%	0.85	0.84	6.29	6.35
T8	NPK @100% + Zn @50% + Fe@100%	0.81	0.92	6.34	6.53
T9	NPK @100% + Zn @100% + Fe @100%	0.91	0.89	6.59	6.84
	F-test	S	S	S	S
	C.D. at 5%	0.265	0.022	0.073	0.081
	S.Ed. (+)	0.012	0.010	0.034	0.038

Table 6. Effect of NPK, Zinc and Iron on Plant height (cm), No. of pods plant⁻¹, No of seeds pod⁻¹ and grain yield (t ha⁻¹) of post-harvest soil

Treatment	Treatment Combination	Plant height (cm)				No of pods plant ⁻¹	No of seeds pod ⁻¹	Grain yield (t ha ⁻¹)
		15 DAS	30 DAS	45 DAS	60 DAS			
T ₁	NPK @0% + Zn @0% + Fe @0%	11.08	22.48	30.41	36.71	17.56	5.17	1.15
T ₂	NPK @0% + Zn @50% + Fe @ 100%	12.36	25.18	33.57	40.78	23.20	7.87	1.78
T ₃	NPK @0% + Zn @100% + Fe @100%	12.55	25.24	33.69	41.14	24.49	8.03	1.81
T ₄	NPK @50% + Zn @0% + Fe @100%	12.80	25.38	33.79	41.27	24.84	7.92	1.87
T ₅	NPK @50% + Zn @50% + Fe @100%	13.83	26.27	37.51	45.25	28.40	8.43	2.06
T ₆	NPK @50% + Zn @100% + Fe @100%	13.55	26.23	37.31	45.14	28.27	8.35	1.91
T ₇	NPK @100% + Zn @0% + Fe @100%	13.42	26.16	35.37	43.34	27.36	8.23	1.96
T ₈	NPK @100% + Zn @50% + Fe@100%	13.10	25.70	35.92	42.21	26.24	8.16	2.03
T ₉	NPK @100% + Zn @100% + Fe@100%	13.92	26.44	38.43	45.53	29.70	8.68	2.09
	F-test	S	S	S	S	S	S	S
	C.D. at 5%	0.156	0.108	0.323	0.849	0.865	0.215	0.808
	S.Ed. (+)	0.073	0.051	0.152	0.400	0.408	0.101	0.381




Fig. 2. Effect of NPK, Zinc and Iron on available Zinc and Iron of Post-harvest soil

4. CONCLUSION

The results of experiment on Effect of NPK, zinc and iron on soil health and yield parameters of greengram was found positively significant on Percentage pore space, Water holding capacity, Percentage organic carbon, Av. Nitrogen, Phosphorus, Potassium, zinc and iron while Bulk density, Particle density and pH and EC was found non-significant. Treatment T₉ [NPK @100% + Zn @100% + Fe @100%] was found most effective in improving Physico-chemical properties of soil. Similarly, the maximum plant height, number of pods plant⁻¹, no of seeds pod⁻¹, Grain yield treatment T₉ [NPK @100% + Zn @100% + Fe @100%]. It is also recorded that treatment T₈ [NPK @100% + Zn @50% + Fe@100%] gave maximum Net Return of ₹ 140975.40 ha⁻¹ with Cost benefit ratio of 1:2.49.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

Words alone cannot convey how much I appreciate my esteemed advisor's advice, assistance, and knowledge. His insightful advice, creative concepts, and diligent demeanour greatly aided me in carrying out my research and

earning my degree. I am appreciative of the Department of Soil Science and Agricultural Chemistry's friends and faculty for their help in providing me with the tools I need to carry out my research project. Additional research is required to validate the results, as they are based on a single season of experimentation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yuvraj D Gavali, Vaishnavi Jaiswala, Ruhidas Ghatak, Kevin Gawali. Effect of varieties and spacing on growth and yield of green gram (*Vigna radiata* L.). International Journal of Environment and Climate Change. 2022;13(5):179-184.
2. Ranpariya VS, Polara KB, Savaliya CM, Modhvadiya VL, Bodar KH. Effect of potassium, zinc and FYM on growth, quality and yield of summer green gram (*Vigna radiata* L.) under medium black calcareous soil. International Journal of Science, Environment and Technology. 2017;;6(4):2242-2247.
3. Rathore KM, Sharma MK, Manoj, Yadav VK, Yadav RK, Meena H, Ghasil BP. Effect of iron, zinc and microbial inoculants on soil health and nutrient uptake of Urdbean (*Vigna mungo* L.) in Vertisols. J. Exp. Agric. Int. 2023, Aug 25;45(10):6-17. [cited 2024 Jun. 2]

- Available:<https://journaljeai.com/index.php/JEAI/article/view/2195>
4. Kamalakannan AS. Assessment of secondary and micronutrients of soil in Coimbatore District, Tamil Nadu, India. *Int. J. Plant Soil Sci.* 2023, Aug 11;35(18):2116-9. [cited 2024 Jun. 2] Available:<https://journalijpss.com/index.php/IJPSS/article/view/3500>
 5. Shankar MA, Sankar GM, Sharma KL, Muniswamappa MV, Rao CS, Chandrika DS. Effect of micronutrient-based integrated use of nutrients on crop productivity, nutrient uptake, and soil fertility in greengram and finger millet sequence under semi-arid tropical conditions. *Communications in Soil Science and Plant Analysis.* 2013, Oct 28;44(19):2771-87.
 6. Tiwari AK, Shivhare AK. Pulses in India retrospect and prospects. Government of India Ministry of Agriculture Department of Agriculture and cooperation Directorate of Pulses Development Vindhyachal Bhavan Bhopal; 2017.
 7. Nair RM, Yang RY, Easdown WJ, Thavarajah D, Thavarajah P, Hughes JDA, Keatinge JDH. Biofortification of mungbean (*Vigna radiata* L.) as a whole food to enhance human health. *Journal of the Science of Food and Agriculture.* 2013; 93(8):1805–1813.
 8. Asif M, Rooney LW, Ali R, Riaz MN. Application and opportunities of pulses in food system. A review: *Critical Reviews in Food Science and Nutrition.* 2013;53: 1168–1179.
 9. Hou D, Yousaf L, Xue Y, Hu J, Wu J, Hu X, Feng N, Shen Q. Mung bean (*Vigna radiata* L.): Bioactive polyphenols, polysaccharides, peptides, and health benefits. *Nutrients.* 2019;11:1238.
 10. Peoples M, Herridge D, Ladha J. Biological Nitrogen fixation: An efficient source of nitrogen for sustainable Agricultural Production. *Plant and Soil Journal.* 2018; 174:3-28.
 11. Dhakal Y, Meena RS, Verma SK, Singh A. Growth, yield and nutrient content of mung bean (*Vigna radiata* L.) in response to INM in eastern Uttar Pradesh, India. *Bangladesh journal of Botany.* 2015; 44(3):479-482.
 12. Mishra PK, Bisht SC, Pooja R, Joshi GK, Singh G, Bisht JK, Bhatt JC. Bio associative effect of cold tolerant pseudomonas spp. and rhizobium leguminosarum-PR 1 on Iron acquisition, nutrient uptake and growth of lentil (*Lens culinaris* L.). *European Journal of Soil Biology.* 2011;47:35-43.
 13. Partha Deb Roy, Lakshman K, Narwal RP, Malik RS, Sushanta Saha. Green gram (*Vigna radiata* L.) Productivity and Grain Quality enrichment through Zinc Fertilization. *Inc. Curr. Microbial. App. Sci.* 2017;6(6):643-648.
 14. Muthuvel P, Udayasoorian C, Natesan R, Ramaswamy PP. Introduction to soil analysis, Tamil Nadu Agricultural University Coimbatore -641002; 1992.
 15. Jackson ML. *Soil chemical analysis* Prentice Hall of India Ltd. New Delhi. 1958;219-221.
 16. Wilcox LV. *Electrical conductivity Am. water work, Association.* 1950;42:775-776.
 17. Walkley A, Black IA. Estimation of soil organic carbon by the chromic acid titration method. *Soil Science;* 1947.
 18. Subbiah BV, Asijja EC. A rapid procedure for estimation of available nitrogen in soil. *Current Science.* 1956;25(8):259-260.
 19. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate (NaHCO₃), U.S.D.A. Circular. 1954;939:1-19.
 20. Toth SJ, Prince AL. Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique *Soil Sci.* 1949; 67:439-445.
 21. Lindsay WL, Norvell WA. Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of America Journal.* 1978;42:421-428.
 22. Chintha HB, Swaroop N, Thomas T, Barthwal A, Harit H, Amjad A. Effect of nitrogen, phosphorus, potassium, sulphur and zinc on soil health parameters of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal.* 2021;10(11):1955-1961.
 23. Hussain M, Banoo M, Sinha BK, Chand. Effect of foliar application of zinc and boron on growth, yield and quality attributes in chickpea (*Cicer arietinum* L.). *Journal of Pharmacognosy and Phytochemistry.* 2022;11(3):270-275.
 24. Prakash Dev Verma, Narendra Swaroop, Yogesh Upadhyay, Akash Swamy, Soman Singh Dhruw. Role of Phosphorus, Zinc and Rhizobium on Physico-Chemical Properties of Soil in Field Pea (*Pisum*

- sativum* L.) var. Rachna. International Journal Current Microbiology and Applied Science. 2017;6(7):4423-4428
25. Banjara GP, Majgahe SK Effect of Biofortification of zinc and iron on yield attributes and yields of chickpea (*Cicer arietinum* L.) through agronomic intervention. The Pharma Innovation Journal. 2019;8(10):45-47.
 26. Tarun Kumar, Tarence Thomas, Narendra Swaroop, Vishal Manohar Dalavi. Influence of zinc and iron on Physico-chemical properties of soil in cluster bean. The Pharma Innovation Journal. 2022;11(8): 1398-1404.
 27. Lokendra Meena, Ram Bharose, Arun Alfred David a, Tarence Thomas, Neha Toppo, Harsh Kumar. Effect of NPK and zinc inorganic fertilizers on soil health, growth and yield of chickpea (*Cicer arietinum* L.). International Journal of Plant & Soil Science. 2023;35(17):155-161.
 28. Sahu A, Swaroop N, David AA, Thomas T. Effect of different levels of NPK and Zinc on Soil health growth and yield of chickpea (*Cicer arietinum* L.) var. PUSA 362. Int. J Curr. Microbial. App. Sci. 2020;9(10):591-597. 20.
 29. Deshlahare H, Banjara GP, Tandon A. Effect of bio fortification of zinc and iron on yields attributes of chickpea (*Cicer arietinum* L.) through agronomic intervention. International Journal of Chemical Studies. 2019;7(4):219-221.
 30. Hemraj Jat, Yadav KK, Singh DP. Effect of soil and foliar application of Zn and Fe on soil fertility after harvest of maize. Biological Forum – An International Journal. 2022;14(1):637-640.
 31. CP Mohammed Nisab, GK Ghosh. Different fractions of zinc and their relationship with physico-chemical properties in red and lateritic soils of Birbhum district, West Bengal. International Journal of Chemical Studies. 2019;7(4):180-185.
 32. Koushik Ghosh, Narendra Swaroop, Tarence Thomas, Jadhav Ravindra. Soil physico-chemical properties as influenced by combined use of npk and zinc at varying levels under black gram (*Vigna mungo* L.) Cultivation in an Inceptisol of Prayagraj, Uttar Pradesh, India. International Journal of Plant & Soil Science. 2022;34(22):1172-1182.
 33. Chethan KV, Arun A David, Terence Thomas, Narendra Swaroop, Smriti Rao, Amreen Hassan. Effect of different levels of N P K and Zn on physicochemical properties of soil growth parameters and yield by pea (*Pisum sativum* L.). Journal of Pharmacognosy and Phytochemistry. 2018;7(3):2212- 2215.
 34. Rohith JR, Arun Alfred David, Tarence Thomas. Effect of different levels of NPK and zinc on physico-chemical properties of soil, growth and yield of pea [*Pisum sativum* L.] Var. Bliss-101. International Journal Current Microbiology and Applied Science. 2020;9(9):3307-3312
 35. Choudhary RS, Yadav BL, Jat R, Jat S. Effect of organic manures and mineral nutrients on growth, yield attributes and yield of Cluster bean [*Cyamopsis tetragonoloba* L.) Taub.]. The Pharma Innovation Journal. 2021;10(7):526-529.
 36. Rajendra Prasad, Yashbir Singh Shivay, Dinesh Kumar. Interactions of zinc with other nutrients in soils and plants - A review. Indian Journal of Fertilisers. 2016; 12(5):16-26.
 37. Fulpagare DD, Patil TD, Thakare RS. Effect of application of iron and zinc on nutrient availability and pearl millet yield in vertisols. International Journal of Chemical Studies. 2018;6:2647-2650.
 38. Sunil Dahiya S, Bhattoo MS, Khedwal RS. Effect of zinc and sulphur on growth, yield and economicsof cluster bean [*Cyamopsis tetragonoloba* (L.) Taub.]. Int. J Curr. Microbiology. App. Sci. 2017;6(11):3744-3751.
 39. Tribhuwana R, Purib AN, Chavana AN. Effect of micronutrient application on growth, yield and quality of green gram growing iron and zinc deficient soils of Renapur Tahsil, Latur. international Journal of Plant & Soil Science. 2024; 36(5):771-779.
 40. Maddila Teja, Abrar Yasin Baba, Venna Gopijagadeeswar Reddy, Thimmisetty Raviteja, Mattepally Vikram Sai, Bada Maheswara Reddy. Performance of sulphur and iron on yield, yield attributes and economics of green gram (*Vigna radiata* L.). Environment and Ecology. 2022;40(4B):2402—2406, October—December 2022 ISSN 0970-0420.
 41. Vinodkumar HV, Channakeshava S, Basavaraja B, Ananathakumar. Effect of soil and foliar application of zinc on growth and yield of green gram (*Vigna radiata* L.).

- International Journal of Current Microbiology and Applied Sciences. ISSN: 2319-7706. 2020;9(4).
42. Boradkar SG, Adsul PB, Shelke MS, Khule YR. Effect of iron and zinc application on soil properties, nutrient uptake and yield of green gram (*Vigna radiata* L.) in Inceptisol. The Pharma Innovation Journal. 2023;12(3):1663-1669.
 43. Balu Ram. Effect of ferrous sulphate on yield parameters of moong bean (*Vigna radiata* L.) in arid region Rajasthan. The Pharma Innovation Journal. 2022;11(5): 1505-1509.
 44. Dogendra kumar sahu, Narendra swaroop, Tarance Thomas, Arun Alfred David, P Smriti Rao. Effect of different doses of sulphur and Zinc with NPK on physicochemical properties of soil and yield attributes of yellow mustard (*Brassica campastris* L.) International Journal of Current Microbiology and Applied Sciences. 2017;6(6).
 45. Fisher RA. Statistical methods and scientific induction. Journal of the royal statistical society series. 1955;17:69- 78.
 46. Likhittakutum, Narendra Swaroop, Tarence Thomas, Akshita Bartwal. Interactive effect of Zinc and Iron on physicochemical properties of soil on Field Pea. Asian Jr. of Microbiol. Biotech. Env. Sc. 2022;26(3): 75-79.
 47. Deb Roy P, Narwal RP, Malik RS, Saha BN, Kumar S. Impact of zinc application on green gram productivity and grain zinc fortification. Journal of Environmental Biology. 2013;54:851-854.
 48. Pulses Revolution from Food to Nutritional Security, Success Report. Crops Division Government of India Ministry of Agriculture & Farmers Welfare; 2017-2018. Available: <http://dpd.gov.in/Retrospect%20and%20Prospect%20From%20Food%20to%20Nutritional%20Security%2018.pdf>.
 49. Raghvani BR, Goyal SN, Patel JC, Malavia DD. Weed competition in mungbean. Indian Journal of Weed Science. 1985; 17(1):18-21.
 50. Singh B, Usha K. Nodulation and symbiotic nitrogen fixation of cowpea genotypes as affected by fertilizer nitrogen. J Plant Nutri. 2003;26(2):463-473.
 51. Jyoti Soni, HS Kushwaha. Effect of foliar spray of zinc and iron on productivity of mung- bean (*Vigna radiata* L. Wilczek). Journal of Pharmacognosy and Phytochemistry. 2020;9(1):108-111.
 52. Gaffar A, Ehsanullah Nadeem E, Sultan HK. Influence of zinc and iron on yield and quality of sugarcane planted under various trench spacing. Pakistan Journal of Agriculture Science. 2011;192:181-189.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/118530>