



# Unveiling the Potential of Nanoparticles in Barley Farming

Tejaswi Akula <sup>a++\*</sup> and Kishore S. M. <sup>b#</sup>

<sup>a</sup> Department of Agronomy, School of Agriculture, Lovely Professional University, Phagwara- 144411, Punjab, India.

<sup>b</sup> Department of Agricultural Entomology, Keladi Shivappa Nayaka University Agricultural and Horticultural Sciences, Shivamogga-577204, Karnataka, India.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i81144>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119938>

Review Article

Received: 15/05/2024

Accepted: 17/07/2024

Published: 20/07/2024

## ABSTRACT

Barley continues to stand out as an important food source for human beings all over the world. It remains significant to manufacturing (industrial applications), especially in the malting and brewing sectors. This contributes to its promotion as a commercial crop through practices such as contract farming in Punjab, Rajasthan, and Haryana. Nanoparticle-based delivery mechanisms are high-end tools for enhancing zinc efficiency in barley crops. These involve coating and stabilizing zinc ions in nanoscale materials such as nanoparticles, nanotubes, or nanocarriers thus improving bioavailability and uptake besides abolishing soil solubility and mobility challenges. In conclusion, nano zinc is a sustainable way of addressing micronutrient deficiencies while promoting barley production under diverse agricultural conditions. This method has huge potential to improve future global food security and agricultural sustainability.

<sup>++</sup>Msc Scholar;

<sup>#</sup>PhD Scholar;

<sup>\*</sup>Corresponding author: E-mail: [akulatejaswi8084@gmail.com](mailto:akulatejaswi8084@gmail.com);

**Cite as:** Akula, Tejaswi, and Kishore S. M. 2024. "Unveiling the Potential of Nanoparticles in Barley Farming". *Journal of Advances in Biology & Biotechnology* 27 (8):315-23. <https://doi.org/10.9734/jabb/2024/v27i81144>.

**Keywords:** *Hordeum vulgare*; seed priming; nano zinc; nutrient uptake; abiotic stress; nanoparticles; agricultural sustainability; food production; nutrition.


## 1. INTRODUCTION

Barley is scientifically known as *Hordeum vulgare* and commonly referred to as “jau” in India. Barley has hardly played a significant role in global production throughout human history. Thus, barley comes fourth among cereal crops currently. This ranks globally at number four with a production of 155 million tons in 2022 after maize, wheat, and rice which make up about 12% of total cereal output. [1]. Barley production thrives in different conditions including the tropical and temperate regions [2-5]. In India, it is majorly cultivated in states such as Rajasthan, Uttar Pradesh, and Madhya Pradesh [6]. The grain has gained a lot of attention for its ability to survive in harsh conditions like drought, excessive heat, and frost [7,8]. Barley is categorized into two genotypes which are two-rowed and six-rowed varieties that have different functions. Since the six-rowed barley is high in protein, it is utilized as animal feed, while the two-row one is traditionally used for malting in Europe, Australia, and south America [9,10]. In agriculture, barley has a wide range of applications. 70% of the global production of barley is used as animal feed, with the remaining supporting beer and whisky production, and for different culinary purposes [11]. In addition, this crop is excellent for pasture, green manure, and as a monoculture, thus reflective of the multifarious value of this crop in farming systems [12]. The malting process, starting from steeping and germination to drying and storage, enhances the utility of barley by rendering it further

processable into malt for brewing and distillation purposes [13]. All byproducts from the malting and brewing industries can be used to manufacture feed and non-feed products, that aid to tackle waste management practices [14]. Further ongoing research is therefore done in investigating the potential of barley in producing modified starches and cereals of high nutritional value, such as high protein or specialized starch compositions that can supply the nutritional fiber, vitamin and protein requirement [15,16]. Thus, barley remains to play a significant part in global agriculture, in food production, and human nutrition by proving its worth across various sectors.

## 2. SIGNIFICANT ADVERSE EFFECTS OF BARLEY PRODUCTION

The poor environment is one of the major challenges to crop production and acts through various means on the growth of barley. The main challenges include drought, salinity, micro and macronutrient deficiency. All these factors lead to a reduction in yield and quality, the effect being more on the six-rowed used for malting because of its low carbohydrate content in the endosperm and high husk ratio, is more accentuated due to inappropriate irrigation and fertilizer application and restricted sowing of malt varieties [17]. However, barley being an important crop in industrial uses, too ensures that with contract farming efforts in states like Punjab, Haryana, rajasthan the crop will certainly come up as a commercial crop [18]. However, the adaptability



**Fig. 1. Significant adverse effects on barley production**

of barley plants to environmental challenges, including drought, has been studied through genetic variation and the development of drought-tolerant genotypes [19,20]. The salinity, which is abundant in semi-arid and arid locations, as well as coastal areas with low-quality irrigation water, serves as another important barrier to barley production, hindering germination and early seedling stages [21,22]. Barley's response to salt stress is driven by ionic responses rather than osmotic stress [23]. To counteract these challenges, potassium nitrate (KNO<sub>3</sub>) is applied to ameliorate oxidative stress and maintain favorable physiological features such as reduced malondialdehyde (MDA) content and altered K<sup>+</sup>/Na<sup>+</sup> concentrations in leaves [24,25] (Fig. 1).

### 3. SEED PRIMING IN BARLEY

Successful seed germination is essential in agriculture and horticulture [26]. Improper germination may result in erratic growth of seedlings and significant financial losses [27]. Seed priming has become more prevalent as a means of rejuvenating seeds and enhancing yield from agriculture [28], notably in commercial applications and seed banks aiming to boost germplasm preservation [29,30]. Primed seeds are hydrated and dried to stimulate metabolic activities, stimulation of enzymes, and antioxidant defenses, resulting in faster and more uniform germination over unprimed seeds [31,32]. There are several priming methods hydropriming, halopriming, Osmo priming, Bio-priming, Nano-priming, Solid matrix priming, and Hormone priming have been developed to enhance seed germination and reduce

environmental stress [33]. The effectiveness of seed priming depends on factors like oxygen availability, priming agent duration, osmotic potential, temperature, light conditions, aeration, and seed condition, which significantly influence its outcomes across different crops [34]. Nano-priming, a novel approach, has shown promising results in enhancing growth, yield, and nutritional quality through its activation of Reactive Oxygen Species (ROS), antioxidant mechanisms, and promotion of nanopore formation for improved water absorption and starch hydrolysis [35,36]. In agriculture, adopting innovative practices like seed priming, including nano-priming, holds the potential to revolutionize crop production to meet future food security and quality challenges [37].

#### 3.1 Overview of Nanoparticles-Based Delivery Systems

Nanoparticle-based delivery systems represent advanced methods for enhancing zinc efficiency in barley plants. These systems utilize nanoscale materials like nanoparticles, nanotubes, or nanocarriers to encapsulate and stabilize zinc ions, improving their bioavailability and uptake by overcoming solubility and mobility barriers in soil [38]. Engineered nanoparticles can release zinc ions gradually, ensuring sustained nutrient supply to plants while protecting against degradation and leaching, thereby optimizing zinc utilization [39,40] (Fig. 2).

Zinc oxide nanoparticles (ZnO NPs) are a type of Zn carriers that help in delivery to roots and improving the uptake as well as translocation with barley and other crops [41].




Fig. 2. Schematic illustration of the wurtzite ZnO structure [42]

This subsequently facilitated the stabilization, dispersion in applied soil, and interaction with plant roots ensuring more absorption of zinc into biomass leading to a boosted crop yield by chitosan or humic acid functionalization [39,38].

### 3.2 Importance of Zinc in Barley Growth and Development

Zinc Importance in Barley Growth and Development Zinc is important for barley growth, it is a co-factor to help enzymes that are necessary during photosynthesis and respiration as well as nutrient synthesis [43]. This mineral controls the level of hormones in a living organism, promoting the elongation of cells in them, their roots, and the flowering of the seeds [44]. The deficit is sure to lead to an inability to synthesize the DNA, divide the cells, or provide the membrane's integrity, decreasing stress resistance and overall vitality [45]. Structurally, zinc holds the integrity of the cell's wall and the stability of the membranes and transportation of ions, which is necessary for taking the diet from the outside and the ability to survive all the dangerous environmental influences [46]. Chlorosis is a deficiency of chlorophyll pigmentation which is essential for photosynthesis, thus without it the plant will not be able to conduct the basic exchange of substances. A lack of zinc is also discovered in a worryingly low tillering and maturation process, which proves that the barley is formed by unhealthy seeds in need of being nourished [43] (Table 1).

### 3.3 Effects of Nano Zinc on Barley Growth Parameters

Nano zinc has a positive effect on the development of barley, contributing to increased quantity and better quality of the yield. This research showed that nano zinc aided in the growth of shoots, increased the amount of plant material, and enhanced photosynthesis [47,48].

Barley treated with Nano zinc has longer roots with more surface area and volume, which helps the plant take in nutrients and water better [47]. It also provides an extension of shoots, expanding the leaf area and enhancing general strength and features of growth in the plants [48]. Nano-sized zinc improved the production of chlorophyll, the rate of photosynthesis, and the absorption of nutrients in barley [49,47]. Additionally, Nano zinc boosts antioxidant enzyme activity, mitigating oxidative stress in barley exposed to drought, salinity, or heavy metals [48,47]. Nano-sized zinc influences the activity of genes that control growth and response to stress, making barley more resistant to living and non-living threats [48,49]. This stress-associated gene control helps the barely plants to strengthen themselves and survive in an unfavorable environment. However, nano-sized zinc is a promising way to increase the yield and hardness of the barley crop (Table 2).

### 3.4 Physiological and Biochemical Responses of Barley to Nano Zinc

Nano zinc improves the growth and quality of barley by enhancing the development of roots and shoots, increasing the amount of biomass produced, and improving the efficiency of photosynthesis. [47,48]. It helps to increase the length and surface area of roots, as well as the uptake of nutrients, while also promoting the growth of shoots and the capacity for tillering [47,48]. Nano zinc also increases the production of chlorophyll, the rate of photosynthesis, and the activity of antioxidant enzymes, which makes barley more resistant to environmental stresses such as drought and saline soils [49,47]. In addition, it regulates the expression of genes to improve tolerance to stress and overall plant health [48,49]. Therefore, the application of nano zinc is beneficial for increasing barley productivity in challenging agricultural conditions (Table 3).

**Table 1. Zinc in barley growth and development**

Aspect	Importance of Zinc in Barley Growth and Development	References
Enzyme Cofactor	➤ Essential for enzyme activity in photosynthesis, respiration, carbohydrate, protein, and nucleic acid synthesis	Marschner, [43]
Hormone Regulation	➤ Involved in synthesis and activation of plant hormones like auxins, regulating cell elongation, division, and differentiation	Alloway, [44]
DNA Synthesis and Cell Integrity	➤ Required in DNA synthesis, cell division, and membrane integrity, thus increasing vigour in plants and their tolerances of stresses.	Cakmak, [45]

Aspect	Importance of Zinc in Barley Growth and Development	References
Structural Integrity	➤ Building block of proteins/complexes involved in cell wall synthesis, stability of membranes, and transport of ions.	Broadley et al., [46]
Symptoms of Deficiency	➤ Barley plants grow with characteristic symptoms of zinc deficiency, developing chlorosis, reduced tillering and delayed maturity.	Marschner, [43]

**Table 2. Nano zinc on barley growth parameters**

Aspect	Effects of Nano Zinc on Barley Plants	References
Growth Parameters	<ul style="list-style-type: none"> <li>➤ Distinctly promotes root and shoot growth.</li> <li>➤ Increases biomass accumulation.</li> <li>➤ Boosts photosynthetic efficiency.</li> </ul>	Shahbaz et al., [47]; Rehman et al., [48]
Morphological Attributes	<ul style="list-style-type: none"> <li>➤ Increases root length, surface area, and volume.</li> <li>➤ Promotes shoot elongation and leaf area expansion.</li> </ul>	Shahbaz et al., [47]; Rehman et al., [48]
Physiological Processes	<ul style="list-style-type: none"> <li>➤ Enhances chlorophyll synthesis and photosynthetic rate.</li> <li>➤ Improves nutrient assimilation</li> <li>➤ Alleviates oxidative stress.</li> </ul>	Ghori et al., [49]; Shahbaz et al., [47]; Rehman et al., [48]

**Table 3. Physiological and biochemical responses of barley to nano zinc**

Aspect	Physiological and Biochemical Responses of Barley to Nano Zinc	References
Antioxidant Enzyme Activities	<ul style="list-style-type: none"> <li>➤ Enhances activities of antioxidant Enzymes (SOD, CAT, POD)</li> <li>➤ Improves scavenging of reactive oxygen species (ROS).</li> <li>➤ Mitigates oxidative stress.</li> </ul>	Ghori et al., [49]; Rehman et al., [48]
Accumulation of Osmolytes and Compatible Solutes	<ul style="list-style-type: none"> <li>➤ Modulates accumulation of osmolytes (proline, sugars, polyamines).</li> <li>➤ Facilitates osmotic adjustment and stress tolerance.</li> </ul>	Shahbaz et al., [47]; Rehman et al., [48]
Influence on Plant Hormones	<ul style="list-style-type: none"> <li>➤ Affects biosynthesis and metabolism of plant hormones (ABA, cytokinins, gibberellins).</li> <li>➤ Alters physiological processes and stress responses.</li> </ul>	Shahbaz et al., [47]
Nutrient Uptake and Assimilation	<ul style="list-style-type: none"> <li>➤ Influences uptake, assimilation, and distribution of essential nutrients (nitrogen, phosphorus, potassium).</li> <li>➤ Enhances nutrient uptake efficiency and Utilization.</li> </ul>	Rehman et al., [48]

#### 4. CHALLENGES AND FUTURE PERSPECTIVES IN NANO ZINC APPLICATION IN BARLEY

Using nano zinc in barley can help it grow better, produce more, and handle stress better.

However, some problems need to be dealt with before it can be used safely and effectively in

farming. One massive concern is how releasing nanoparticles into the environment could harm soil and water ecosystems. There's a chance that these particles could stick around for a long time and build up in non-target animals, which could be toxic for them [50]. To mitigate these risks, rigorous risk assessments and regulatory frameworks are essential to ensure sustainable Nano zinc use.

Understanding the fate of Nano zinc particles in soil-plant systems remains limited, impacting predictions of their effects on soil health, microbial communities, and nutrient cycling [51]. Interactions with soil components like organic matter and minerals affect mobility and bioavailability, necessitating further research to clarify these mechanisms and their implications for biogeochemical processes. Variability in Nano zinc formulations, including particle size, surface chemistry, and coatings, poses another challenge that can influence effectiveness and behaviour in agricultural contexts [47]. Standardized protocols for synthesizing and characterizing Nano zinc are crucial to ensure consistency across studies and facilitate meaningful comparisons. Despite these challenges, Nano zinc shows promise in addressing barley's micronutrient deficiencies and improving crop productivity sustainably. Future research should focus on optimizing formulations, dosage, and application methods to maximize benefits while minimizing environmental risks [47]. Integrating Nano zinc with precision farming, biofortification, and nutrient management strategies could further enhance its efficacy across diverse agroecosystems [48].

## 5. CONCLUSION

In conclusion, nanoparticle-based Nano zinc application in barley cultivation offers a promising approach to enhance crop productivity, stress tolerance, and nutritional quality. Research has shown that Nano zinc positively impacts various aspects of barley growth, including root and shoot development, photosynthetic efficiency, and antioxidant defenses. By improving nutrient uptake, reducing oxidative stress, and regulating hormone levels, Nano zinc enhances barley's ability to withstand environmental stresses and achieve higher yields. Despite these benefits, challenges such as environmental risks, variability in nanoparticle formulations, and gaps in understanding Nano zinc's behavior in soil-plant systems need addressing for safe and effective agricultural use. Moving forward, further research should prioritize optimizing Nano zinc formulations, dosage, and application methods. Integrating Nano zinc with other agricultural practices could maximize its benefits while minimizing potential drawbacks.

Overall, Nano zinc represents a promising and sustainable solution for combating micronutrient deficiencies and enhancing barley production in

diverse agricultural settings. This approach holds significant potential to contribute to global food security and agricultural sustainability in the future.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## ACKNOWLEDGEMENT

The authors express their sincere gratitude towards their parents for providing them with constant support and encouragement.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Verma RPS, Lal C, Malik R, Kharub AS, Kumar L, Kumar D. Barley improvement: Current status and future prospects in changing scenario. *New Horizons in Wheat and Barley Research: Global Trends, Breeding and Quality Enhancement*. 2022;93-134.
2. Verma, Sandeep Kumar, Rana NS, Aparna Verma, Shalini Roy, Aditya Shukla, Ajay Kumar. Effect of foliar application of nano and non-nano fertilizers on growth and nutrient use efficiency of Indian Mustard [*Brassica juncea* (L.) Czern and Coss]. *Journal of Scientific Research and Reports*. 2024;30(5):868-78. Available:<https://doi.org/10.9734/jsrr/2024/v30i52004>
3. Abhilash, Akula, Abhinav Dayal, Neha Thomas, Abhishek Sharan, Vadlakonda Vipul. Effect of Zinc Oxide (ZnO) Nanoparticles on the Storability of Onion (*Allium cepa* L.) Seeds under Ambient Condition. *International Journal of Plant & Soil Science*. 2023;35(22):455-68. Available:<https://doi.org/10.9734/ijpss/2023/v35i224154>
4. Faizan M, Hayat S, Pichtel J. Effects of zinc oxide nanoparticles on crop plants: A perspective analysis. *Sustainable Agriculture Reviews* 41: Nanotechnology

- for Plant Growth and Development. 2020;83-99.  
Available:[https://link.springer.com/chapter/10.1007/978-3-030-33996-8\\_4](https://link.springer.com/chapter/10.1007/978-3-030-33996-8_4)
5. Rossi L, Fedenia LN, Sharifan H, Ma X, Lombardini L. Effects of foliar application of zinc sulfate and zinc nanoparticles in coffee (*Coffea arabica* L.) plants. Plant Physiology and Biochemistry. 2019 Feb 1;135:160-6.  
Available:<https://www.sciencedirect.com/science/article/abs/pii/S0981942818305448>
  6. Shekhawat PS, Singh SP, Bishnoi S. Integrated management of barley diseases: Current status and future research priorities. In New Horizons in Wheat and Barley Research: Crop Protection and Resource Management. Singapore: Springer Nature Singapore. 2022;215-26).
  7. Newton AC, Flavell AJ, George TS, Leat P, Mullholland B, Ramsay L, Bingham IJ. Crops that feed the world 4. Barley: A resilient crop? Strengths and weaknesses in the context of food security. Food Security. 2011;3:141-178.
  8. Wang J, Vanga SK, Saxena R, Orsat V, Raghavan V. Effect of climate change on the yield of cereal crops: A review. Climate. 2018;6(2):41.
  9. Öztürk I, Tülek B, Remzi AVCI, Kahraman T, Tülek A. Comparison of two and six-rowed barley (*Hordeum vulgare* L.) genotypes under rainfed conditions for yield, quality and biotic stress tolerance. Ekin Journal of Crop Breeding and Genetics. 2023;9(2):98-105.
  10. Horsley RD, Franckowiak JD, Schwarz PB. Barley. Cereals. 2009;227-250.
  11. Edney MJ. Barley. In Cereal grain quality. Dordrecht: Springer Netherlands. 1996; 113-151.
  12. Steinbach J. Alternatives to crop residues as feed resources in mixed farming systems. Crop Residues in Sustainable Mixed Crop/Livestock Farming Systems. 1997;10:197.
  13. Patel A, Devraja HC, Sharma P, Singh RRB. Food Technology II; 2019.
  14. Karlovic A, Juric A, Coric N, Habschied K, Krstanovic V, Mastanjevic K. By-products in the malting and brewing industries—re-usage possibilities. 2020;6(3):82.
  15. Jadhav SJ, Lutz SE, Ghorpade VM, Salunkhe DK. Barley: Chemistry and value-added processing. Critical Reviews in Food Science.1998;38(2):123-171.
  16. Zohoori FV. Nutrition and diet. The Impact of Nutrition and Diet on Oral Health. 2020;28:1-13.
  17. Hajiboland R. Effect of micronutrient deficiencies on plants stress responses. Abiotic Stress Responses in Plants: Metabolism, Productivity and Sustainability. 2012;283-329.
  18. Kumar V, Khippal A, Singh J, Selvakumar R, Malik R, Kumar D, Sharma I. Barley research in India: Retrospect and prospects. Journal of Wheat Research. 2014;6(1):1-20.
  19. Bukhari SAH, Peerzada AM, Javed MH, Dawood M, Hussain N, Ahmad S. Growth and development dynamics in agronomic crops under environmental stress. Agronomic Crops: Production Technologies. 2019;1:83-114.
  20. Serraj R, McNally KL, Slamet-Loedin I, Kohli A, Haefele SM, Atlin G, Kumar A. Drought resistance improvement in rice: An integrated genetic and resource management strategy. Plant Production Science. 2011;14(1):1-14.
  21. Pulido-Bosch A, Rigol-Sanchez JP, Vallejos A, Andreu JM, Ceron JC, Molina-Sanchez L, Sola F. Impacts of agricultural irrigation on groundwater salinity. Environmental Earth Sciences. 2018;77:1-14.
  22. Fan Y, Zhu M, Shabala S, Li CD, Johnson P, Zhou MX. Antioxidant activity in salt-stressed barley leaves: Evaluating time-and age-dependence and suitability for the use as a biochemical marker in breeding programs. Journal of Agronomy and Crop Science. 2014;200(4): 261-272.
  23. Mahlooji M, Seyed Sharifi R, Razmjoo J, Sabzalian MR, Sedghi M. Effect of salt stress on photosynthesis and physiological parameters of three contrasting barley genotypes. Photosynthetica. 2018;56:549-556.
  24. Ahmad S, Wang GY, Muhammad I, Zeeshan M, Zhou XB. Melatonin and KNO<sub>3</sub> application improve growth, physiological and biochemical characteristics of maize seedlings under waterlogging stress conditions. Biology. 2022;11(1):99.
  25. Sofy MR, Elhawat N, Alshaal T. Glycine betaine counters salinity stress by maintaining high K<sup>+</sup>/Na<sup>+</sup> ratio and antioxidant defense via limiting Na<sup>+</sup> uptake in common bean (*Phaseolus vulgaris* L.).

- Ecotoxicology and Environmental Safety. 2022;200:110732.
26. Kulkarni MG, Light ME, Van Staden J. Plant-derived smoke: Old technology with possibilities for economic applications in agriculture and horticulture. *South African Journal of Botany*. 2011;77(4):972-979.
  27. Bisbis MB, Gruda N, Blanke M. Potential impacts of climate change on vegetable production and product quality—A review. *Journal of Cleaner Production*. 2018;170:1602-1620.
  28. Marthandan V, Geetha R, Kumutha K, Renganathan VG, Karthikeyan A, Ramalingam J. Seed priming: A feasible strategy to enhance drought tolerance in crop plants. *International Journal of Molecular Sciences*. 2020;21(21):8258.
  29. Farooq M, Usman M, Nadeem F, ur Rehman H, Wahid A, Basra SM, Siddique KH. Seed priming in field crops: Potential benefits, adoption and challenges. *Crop and Pasture Science*. 2019;70(9):731-771.
  30. Sershen D, Hay FR. New technologies to improve the ex situ conservation of plant genetic resources; 2021.
  31. Jisha KC, Vijayakumari K, Puthur JT. Seed priming for abiotic stress tolerance: An overview. *Acta Physiologiae Plantarum*. 2013;35, 1381-1396.
  32. Paul S, Dey S, Kundu R. Seed priming: an emerging tool towards sustainable agriculture. *Plant Growth Regulation*. 2022; 97(2):215-234.
  33. Pawar VA, Laware SL. Seed priming a critical review. *Int. J. Sci. Res. Biol. Sci*. 2018;5(5):94-101.
  34. Lutts S, Benincasa P, Wojtyla L, Kubala S, Pace R, Lechowska K, Garnczarska M. Seed priming: New comprehensive approaches for an old empirical technique. *New Challenges in Seed Biology-basic and Translational Research Driving Seed Technology*. 2016;46.
  35. do Espirito Santo Pereira A, Caixeta Oliveira H, Fernandes Fraceto L, Santaella C. Nanotechnology potential in seed priming for sustainable agriculture. *Nanomaterials*. 2021;11(2):267.
  36. Nile SH, Thiruvengadam M, Wang Y, Samynathan R, Shariati MA, Rebezov M, Kai G. Nano-priming as emerging seed priming technology for sustainable agriculture—recent developments and future perspectives. *Journal of Nanobiotechnology*. 2022;20(1): 254.
  37. Beddington J. Food security: Contributions from science to a new and greener revolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2010;365(1537):61-71.
  38. Khan, Tariq, et al. Plant-based gold nanoparticles; a comprehensive review of the decade-long research on synthesis, mechanistic aspects and diverse applications. *Advances in Colloid and Interface Science*. 2019;272:102017.
  39. Rizwan M, Ali S, Adrees M, Rizvi H, Rehman MZ, Hannan F, Al-Wabel MI. Zinc and iron oxide nanoparticles improved the plant growth and reduced the oxidative stress and cadmium concentration in wheat. *Chemosphere*. 2019;214:269-277.
  40. Zhu H, Han J, Xiao JQ, Jin Y, Uchimiya M. Immobilization of metal oxide nanoparticles on montmorillonite modified with cysteine for enhanced adsorption of copper (II) and zinc (II). *Journal of Environmental Management*. 2018;217:568-574.
  41. Eichert T, Kurtz A, Steiner U. Gold, platinum, and titanium dioxide nanoparticles do not affect soil zinc availability to wheat. *Journal of Plant Nutrition and Soil Science*. 2011;174(4): 597-599.
  42. Ding, Meng, Zhen Guo, Lianqun Zhou, Xuan Fang, Lili Zhang, Leyong Zeng, Lina Xie, and Hongbin Zhao. One-Dimensional Zinc Oxide Nanomaterials for Application in High-Performance Advanced Optoelectronic Devices *Crystals* 8. 2018;5: 223. Available:<https://doi.org/10.3390/cryst8050223>
  43. Marschner, Horst, ed. *Marschner's mineral nutrition of higher plants*. Academic Press, Crops; 2012.
  44. Alloway BJ. *Zinc in soils and crop nutrition* (2nd ed.). International Zinc Association, Brussels, Belgium; 2008.
  45. Cakmak, Ismail. Enrichment of cereal grains with zinc: Agronomic or genetic biofortification?. *Plant and Soil*. 2008; 1-17.
  46. Broadley MR, White PJ, Hammond JP, Zelko I, Lux. Zinc in plants. *New Phytologist*. 2007;173(4):677-702.
  47. Shahbaz M, Ashraf MY, Al-Qurainy F, Harris PJC. Nanoparticles impact growth



- and physiological activities of crop plants: A focus on wheat, maize, and barley. *Molecules*. 2020;25(13):2924.
48. Rehman MZU, Khan S, Brusseau ML, Shah MT, Din IU. Effect of nano zinc oxide on growth, photosynthetic pigments, and antioxidant enzyme activities of oat (*Avena sativa* L.) and barley (*Hordeum vulgare* L.) under different soil moisture levels. *Environmental Science and Pollution Research*. 2019;26(23):23933-23944.
49. Ghori NH, Ghori T, Hayat MQ, Imadi SR, Gul A, Altay V, Ozturk M. Heavy metal stress and responses in plants. *International Journal of Environmental Science and Technology*. 2019;16:1807-1828.
50. Judy JD, Unrine JM, Bertsch PM, Hunyadi SE. Exposing engineered nanoparticles to contaminated environments: Environmental fate, transformation, and toxicity of nanoparticles. In *Engineered nanoparticles and the environment: Biophysicochemical Processes and Toxicity*. John Wiley & Sons. 2015;213-238.
51. Dimkpa CO, McLean JE, Latta DE. Fate of CuO and ZnO nano-and microparticles in the plant environment. *Environmental Science & Technology*. 2015;49(17): 10706-10713.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/119938>