



# A Comprehensive Review of Advances in Semiochemical Exploitation for Insect Pest Management

**Shashank D. U. <sup>a\*</sup>, Sujatha G S <sup>b</sup>,  
Katta Subramanya Sai Teja <sup>a</sup>, Hema A P <sup>c</sup>, Ritu Mishra <sup>d</sup>,  
Abinaya S <sup>e</sup>, K. Shewta <sup>f</sup> and Gadde Anil Kumar <sup>g</sup>**

<sup>a</sup> Department of Entomology, Post Graduate College of Agriculture, Dr. Rajendra Prasad Central Agricultural University, Pusa, Samastipur, 848 125, Bihar, India.

<sup>b</sup> Division of Entomology, ICAR-Indian Agricultural Research Institute, New Delhi- 110012, India.

<sup>c</sup> Department of Entomology, University of Agricultural Sciences, Bengaluru- 560065, India.

<sup>d</sup> Department of Entomology, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar-263145, India.

<sup>e</sup> Department of Entomology, University of Agricultural Sciences, Raichur- 584104, India.

<sup>f</sup> Department of Entomology, Tamil Nadu Agricultural University, 641003, India.

<sup>g</sup> Division of Entomology, Acharya N. G. Ranga Agricultural University, Andhra Pradesh- 522101, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/jabb/2024/v27i91357>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121986>

**Review Article**

**Received: 22/06/2024**

**Accepted: 01/09/2024**

**Published: 05/09/2024**

\*Corresponding author: E-mail: shankydu98@gmail.com;

**Cite as:** D. U., Shashank, Sujatha G S, Katta Subramanya Sai Teja, Hema A P, Ritu Mishra, Abinaya S, K. Shewta, and Gadde Anil Kumar. 2024. "A Comprehensive Review of Advances in Semiochemical Exploitation for Insect Pest Management". *Journal of Advances in Biology & Biotechnology* 27 (9):845-63. <https://doi.org/10.9734/jabb/2024/v27i91357>.

## ABSTRACT

In recent years, semiochemical-based pest management strategies have gained significant attention as they offer sustainable alternatives to conventional pesticides. Integrating semiochemicals into pest management strategies presents innovative approaches to addressing agricultural challenges. One promising method involves combining pheromones with entomopathogenic fungi, utilizing a "lure and infect" technique that attracts pests to fungal pathogens, enhancing control efficacy. Another advancement is the auto-dissemination approach, which promotes the spread of microbial pathogens within insect populations, effectively targeting pests like the fall armyworm. Additionally, the induction of plant defenses through "plant vaccination" by zoo phytophagous predators offers a novel way to enhance plant resistance against herbivores. Research into the production of insect pheromones in plants further supports sustainable pest management by disrupting pest mating behaviors. Electroantennography has emerged as a valuable tool for understanding insect olfaction, aiding in the identification of effective semiochemicals. The push-pull strategy employs plant semiochemicals to manipulate pest behavior, while pheromone dispensers provide efficient and long-lasting applications of these compounds. Collectively, these advancements highlight the potential of semiochemicals in revolutionizing pest management practices, aligning with the increasing demand for sustainable agricultural solutions. Continued research and innovation in these areas are crucial for optimizing the use of semiochemicals, ultimately contributing to more effective and environmentally friendly pest control methods.

*Keywords: Auto-dissemination; electroantennography; pheromone dispensers; plant vaccination; push-pull strategy; semiochemicals; trapping techniques.*

## 1. INTRODUCTION

The 21st century is a time of great challenges for global agriculture, with the world population expected to reach about 10 billion by 2050 [111]. This demographic shift requires a significant increase in crop productivity; however, at the same time, there is growing pressure to reduce the use of conventional chemical pesticides due to environmental and health concerns [92]. Consumers are increasingly demanding safe, healthy food with minimal chemical residues, driving the need for sustainable agricultural practices [96]. Innovative solutions are required to address these complex challenges to enhance crop yield and quality while managing pest resistance and minimizing environmental impact. In this context, semiochemicals have emerged as a promising alternative to conventional pesticides, offering a more targeted and environmentally friendly approach to pest management [90]. Semiochemicals are chemicals that mediate interactions between organisms [1,5,78], derived from the Greek word "semeion" meaning sign or signal. These compounds can be classified into two main categories: pheromones that facilitate communication within the species and allelochemicals that mediate interactions between different species [120]. The application of semiochemicals in agriculture aligns with the

principles of Integrated Pest Management (IPM), providing effective tools for pest control while reducing reliance on broad-spectrum insecticides [121,122]. While extensive research has been conducted on lepidopteran pheromones, recent studies have expanded to non-lepidopteran groups including Hymenoptera, Isoptera, and Hemiptera [127]. This broadening scope of research reflects the growing recognition of semiochemicals' potential across diverse pest species.

Advances in molecular research have greatly improved our understanding of the mechanisms underlying semiochemical action. Studies on pheromone receptors (PRs) have shed light on how insects recognize and decode pheromone signals, providing valuable insights for the development of more effective pest management strategies [108]. This molecular-level understanding is crucial for optimizing the use of semiochemicals in agricultural applications. In addition to basic research, practical innovations in semiochemical deployment are gaining attention. The exploration of meso- and mega-pheromone dispensers represents a potential breakthrough in pheromone-based pest control methods [73,75]. These novel dispensing systems offer the promise of more efficient and long-lasting semiochemical applications, potentially enhancing the efficacy and economic

viability of pheromone-based pest management strategies. Semiochemicals play diverse roles in insect ecology, influencing behaviors such as mate location, aggregation, host finding, and oviposition site selection [2,8]. By manipulating these natural chemical signals, it is possible to disrupt pest behaviours, attract pests to traps or repel them from crops offering multiple avenues for pest control [1,15,17]. Furthermore, semiochemicals can be used to influence the behavior of natural enemies of pests offering opportunities for biological control within IPM frameworks [1]. As we continue to face the dual challenges of increasing agricultural productivity and environmental sustainability, exploiting semiochemicals for insect pest management offers a promising way forward. This review article aims at exploring recent advances in semiochemical research and application highlighting their potential to revolutionize pest management practices in the 21<sup>st</sup> century and beyond.

## 2. SEMIOCHEMICALS IN IPM: MARKET TRENDS AND TYPES

Semiochemicals are chemical substances that mediate communication between organisms,

playing a crucial role in insect behaviour and ecology. These compounds convey information about various aspects of the environment, such as mate location, territory marking, danger signaling, and coordination of group activities [18,20]. The global semiochemicals market was valued at \$4.74 billion in 2022 and is expected to grow at a CAGR of 15.84 per cent from 2023 to 2030 to reach \$15.36 billion by 2030 [105]. Insect sex pheromones currently dominate the semiochemicals market with a share of 69 per cent, followed by aggregation pheromones at 25 per cent [97,98]. These chemicals play a crucial role in managing lepidopteran pests which account for approximately 82 per cent of their use while dipteran pests constitute about 7 per cent [97]. The Asia pacific region is expected to witness the fastest growth in the semiochemicals market due to factors such as increased government support, adoption of sustainable agricultural practices and rise in pest-related challenges. Semiochemicals also play an important role in pre-border biosecurity providing effective tools for the detection and eradication of invasive species thereby contributing significantly towards global agricultural sustainability and pest management [97].

**Table 1. Comprehensive Classification of Semiochemicals and Their Functions**

Type	Subtype	Function	Example	Target Organism
Pheromones	Aggregation	Attract both sexes to food sources or habitats	3-methylbut-3-en-1-ol	Beetles
	Alarm	Alert conspecifics to threats	(E)- $\beta$ -farnesene	Aphids
	Sex	Mediate mate attraction	Bombykol	Silk moths
	Trail	Guide social insects to food sources	(Z)-9-hexadecenal	Ants
	Oviposition-deterrent	Discourage egg-laying in occupied resources	Methyl eugenol	Fruit flies
	Home recognition	Enable colony members to identify nest	Cuticular hydrocarbons	Social insects
	Royal	Enable recognition of reproductive individuals	Heneicosane	Termites
Allelochemicals	Allomones	Benefit emitter (e.g., defense)	Nicotine	Tobacco plants
	Kairomones	Benefit receiver (e.g., prey location)	Bark beetle aggregation pheromones	Predatory beetles
	Synomones	Benefit both emitter and receiver	Floral scents	Plants and pollinators
	Antimones	Maladaptive for both emitter and receiver	Geosmin	Various organisms
	Apneumones	Emitted by non-living sources	Green leaf volatiles	Plants (post-damage)

Semiochemicals can be broadly classified into two main categories: pheromones that facilitate intraspecific communication; and allelochemicals that mediate interspecific interactions (Table 1). Pheromones are species-specific chemical signals that enable communication between individuals of the same species. These compounds can trigger behavioural or physiological changes in the recipient [52]. Several types of pheromones have been identified based on their functions [21,26]. Aggregation pheromones attract individuals of both sexes to food sources or reproductive habitats. For example, the hemiterpene 3-methylbut-3-en-1-ol serves as an aggregation pheromone for certain beetle pests [12]. Alarm pheromones alert conspecifics to potential threats. Sesquiterpenes like (E)- $\beta$ -farnesene and germacrene A are common components of aphid alarm pheromones [113, 7]. Oviposition-deterrent pheromones discourage females from laying eggs in already occupied resources. This behavior has been observed in fruit flies like *Rhagoletis pomonella* (Walsh) [93]. Home recognition pheromones such as queen pheromones in social insects enable colony members to identify their nest and queen [66]. Sex pheromones mediate mate attraction and are primarily produced by females to attract males. The first characterization of a sex pheromone was reported in the silk moth *Bombyx mori* [14]. Trail pheromones guide social insects to food sources exhibiting both recruitment and orientation effects [65]. Recruitment pheromones induce nestmates to leave the nest and migrate to work sites. These are often discharged from specialized exocrine glands [66]. Royal pheromones such as hydrocarbon heneicosane in termites enable workers to recognize reproductive individuals and maintain caste division [31,35].

Allelochemicals facilitate communication between different species and can be further categorized based on benefits conferred to the emitter and receiver [26]. Allomones benefit the emitter, often serving as defense mechanisms against predators or herbivores. For example, nicotine produced by tobacco plants acts as a deterrent to herbivorous insects [117]. Kairomones benefit the receiver such as predators using prey pheromones to locate their targets [24]. One example is the orientation of checkered beetles to bark beetle aggregation pheromones [91]. Synomones are advantageous to both the emitter and receiver. Floral scents that attract pollinators and herbivore-induced

plant volatiles that recruit natural enemies of insect pests are examples of synomones [109,110]. Antimones are maladaptive for both the emitter and receiver, often eliciting repellent responses in interspecific interactions [117]. Apneumones are emitted by non-living sources and elicit responses in receiving organisms that may be beneficial to some species but harmful to others [13].

The understanding of semiochemicals has led to their extensive use in IPM strategies. Pheromones, in particular, have found widespread applications. Pheromone traps are used to monitor and detect pest populations, allowing for timely interventions [37]. Mass trapping through large-scale deployment of pheromone traps can significantly reduce pest populations. Mating disruption, which involves the release of synthetic sex pheromones, can interfere with mate location, reducing pest reproduction. The attract-and-kill strategy combines pheromones with insecticides to lure and eliminate pests [95]. Other semiochemicals also play important roles in pest management. Attractants and repellents derived from various allelochemicals can be used to manipulate pest behaviour. Kairomones are employed in stimulo-deterrent diversion strategies, where pests are lured away from crops and toward trap crops or other control measures [95].

### **3. SEMIOCHEMICALS IN INSECT-PLANT INTERACTIONS: MECHANISMS, EFFECTS, AND APPLICATIONS IN PEST MANAGEMENT**

Insects are affected by semiochemicals that mediate different behaviors and physiological responses. Many active compounds have been isolated and identified from plant materials through phytochemical investigations, which have shown their various effects on insect pests [39,40,41,42,101]. These plant volatiles serve multiple functions for herbivorous insects such as food source identification, mates' location, oviposition sites detection and hibernation locations [26]. Insects detect these compounds by stimulating chemoreceptor cells of taste sensilla present on antennae, tarsi and mouthparts [74]. The response of insects to plant volatiles can vary significantly from attraction to repulsion depending on the insect species and its adaptation to the host plant [26]. It is important to note that the classification of plant volatiles as attractants or repellents is not standardized

because insect behavioral responses may fluctuate with concentration of these compounds [94]. Some insects have evolved to utilize host plant compounds as precursors for sex pheromones or as sex pheromones themselves. For example, male orchid bees collect terpenoid mixtures from orchids which they use as aggregation pheromones during mating [25]. Additionally, pyrrolizidine alkaloids are used by various insect species including moths, butterflies, grasshoppers, beetles and aphids etc. as feeding deterrents against their natural enemies [77].


The efficacy of semiochemicals in insect control strategies depends on several factors such as release rate and trap design. The release rate of semiochemicals is critical for trapping success because high levels of release may not necessarily result in increased insect captures and can even become repellent near of the trap. For example, studies on the red flour beetle, *Tribolium castaneum*, have shown that high release rates of pheromones were neither attractive nor repellent to the beetles while older traps with lower release rates were more effective [46, 89]. Trap design is another crucial factor affecting the efficacy of semiochemical-based control strategies. Different aspects of trap design such as shape, size, height, alignment with wind direction, position and timing can significantly affect catching efficacy [26]. Common types of traps include sticky traps, water traps, and inverted cone traps each having specific advantages for different monitoring or control purposes. Combining chemical and visual stimuli in trap design has been shown to enhance insect responses to lures [106].

#### 4. ADVANCES IN THE USE OF SEMIOCHEMICALS IN INSECT PEST MANAGEMENT

##### **New Trapping Technique by Combining Pheromones with Entomopathogenic Fungi:**

In recent years, the use of semiochemicals combined with entomopathogenic fungi has emerged as a promising strategy for pest management in agriculture. This method is often referred to as “lure and infect” and it utilizes natural dispersal mechanisms of pathogens while increasing their effectiveness through targeted application [114]. Entomopathogens are naturally dispersed through various means including aerial movement of spores, parasitoids, predators, and

insect vectors [70]. However, the “lure and infect” strategy uses semiochemicals to attract pest insects to devices containing entomopathogenic fungi [6]. This allows target pests to be contaminated with the pathogen which they then spread to conspecifics leading to epizootic outbreaks in the pest population [57-59]. The efficacy of this approach has been demonstrated in different pest management scenarios. Mfuti et al. (2017) investigated the combined use of *Metarhizium anisopliae* and the thrips attractant Lurem-TR for managing bean flower thrips on cowpeas. Their field experiments compared spot and cover spray applications where spot application lure and infect were most effective. Thrips density was reduced by this approach from  $67.4 \pm 10.30$  thrips per plant in control to  $30.5 \pm 6.0$  per plant. Importantly, this method required only 81.2 per cent less inoculum than cover sprays making it cost-effective for small-scale farmers. Hajjar et al. (2015) studied on red palm weevil using pheromone traps combined with the entomopathogenic fungus *Beauveria bassiana*. They designed a pheromone trap with internal surfaces coated with *B. bassiana*-treated fabric, which effectively attracted and infected both male and female weevils for up to 13 days after application. The study demonstrated high efficacy of *B. bassiana* at  $1 \times 10^9$  spores per ml, with complete mortality observed even 13 days post-treatment. Kabaluk et al. (2015) did research on click beetles focused on an “attract and kill” strategy that combined a granular formulation of female sex pheromone with the entomopathogenic fungus *Metarhizium brunneum*. Banded applications of pheromone granules with *M. brunneum* conidia reduced beetle recapture by 98.2 per cent compared to *M. brunneum* alone. The pheromone bands significantly increased beetle attraction, allowing them to acquire lethal doses of fungal conidia rapidly. For example, successful applications have been reported for bark beetles (*Ips typographus*), weevils (*Cylas formicarius* and *Cosmopolites sordidus*), moths (*Plutella xylostella*), stink bugs (*Plautia crossota stali*), thrips (*Megalurothrips sjostedti*), and aphids (*Phorodon humuli*) [47,71,114]. However, despite these successes there are still challenges in optimizing the lure and infect strategy. In some cases, sex-specific semiochemicals that attract only one sex may limit its effectiveness [121]. Additionally, applying this approach to soil-dwelling insects presents unique difficulties although promising developments have been made [61,87].



**Fig. 1. Lure and infect strategy for pest management**

For instance, Agriculture and Agri-Food Canada has developed prototype granules that combine *Metarhizium brunneum* with pheromone compounds, which have potential attract *Agriotes cutworms* [49]. To make the “lure and infect” method more effective, it is necessary to consider carefully the physical separation between semiochemicals and entomopathogenic fungi in the delivery system [83]. This spatial arrangement is critical for achieving optimal autoinoculation and ensuring successful infection of target pests. As research in this field advances, further refinements of the “lure and infect” strategy are likely to increase its applicability across a wider range of pest species and agricultural settings. The integration of entomopathogenic fungi with semiochemicals offers a promising avenue for the future development of more sustainable and targeted pest management solutions.

**Auto-dissemination approach:** Auto-dissemination is an innovative approach aimed at the targeted application of microbes that promotes transmission of infective pathogen propagules within and between insect populations [115]. This technique has shown promise in managing *Spodoptera frugiperda* (fall armyworm) adults using certain entomopathogenic fungi such as *Beauveria bassiana* or *Metarhizium anisopliae* isolates. Cárcamo et al. (2020) conducted research and showed that *M. anisopliae* Ma-San Rafael-2 isolate combined with synthetic sex pheromone not only attracted but also killed male moths thus indicating its potential for fall armyworm control. Therefore, the auto-dissemination approach is a promising tool for improving the efficacy of microbial control agents in managing fall armyworm populations.

**Plant vaccination by zoo phytophagous predators beyond biocontrol:** Plant vaccination

by zoo phytophagous predators is a promising approach to enhancing plant resistance against insect pests and diseases [85]. These unique predators, which feed on both insects and plants, have been shown to induce plant defenses through their phytophagy, resulting in complex plant responses that can enhance overall plant protection [76,34,81]. The induced plant responses typically manifest as the emission of volatile organic compounds (VOCs) or the transcription of defense-related genes [11,51, 119]. These responses can exert plant protection effects through various mechanisms. For instance, they may repel arthropod pests directly or attract their natural enemies, thereby indirectly reducing pest populations [22, 53]. Additionally, these induced defenses can activate defense mechanisms in neighbouring plants, preparing them against future attacks in a phenomenon known as plant priming [63, 64, 16]. Among the various zoo phytophagous predators studied, several groups have received extensive attention for their ability to induce plant defences. Pappas et al. (2020) explored innovative biological and molecular approaches for enhancing plant defence against pests and pathogens in agriculture. They highlighted the use of beneficial organisms such as zoophytophagous predators as “plant vaccination agents”. This concept may involve the induction of plant defences including production of volatile organic compounds that act as semiochemicals in interactions between plants and insects. De Puyseleyn (2011, 2014) examined interactions between two economically important zoophytophagous predators, *Nesidiocoris tenuis* and *Orius laevigatus*, and their host plants. The study revealed that both predators activate a jasmonic acid (JA)-mediated wound response in their host plants. Importantly, *N. tenuis* was found to induce the emission of volatile compounds like p-cymene, which can have repellent or toxic effects on herbivores. This finding directly relates to semiochemical-

mediated interactions, as plant-derived volatiles can act as important signalling molecules in tritrophic systems [116]. In a review of zoophytophagous mirid bugs as biological control agents, Pérez-Hedo et al. (2020) discussed the potential of harnessing mirid-induced plant defences for pest management. While not explicitly focused on semiochemicals, this approach likely involves changes in plant volatile profiles that could affect pest behaviour and plant-insect interactions [84]. Predatory mites for example have been shown to elicit strong defensive responses in various plant species [102,19]. Anthocorid bugs particularly those belonging to the genus *Orius* have also been demonstrated to induce significant plant defences [10,86]. Pappas et al. (2015) investigated the effects of the omnivorous predator *Macrolophus pygmaeus* on tomato plant resistance against two herbivore species. They found that prior exposure of tomato plants to *M. pygmaeus* reduced the performance of the two-spotted spider mite (*Tetranychus urticae*) but not the greenhouse whitefly (*Trialeurodes vaporariorum*). This induced resistance was associated with increased accumulation of proteinase inhibitor transcripts and activity, both locally and systemically in the plant. While not explicitly studying semiochemicals, this research demonstrates that zoophytophagous predators can induce plant defence responses that may involve changes in plant volatile emissions. The growing body of research on zoophytophagous predator-induced plant defences highlights the potential of these organisms as biological control agents and as tools for enhancing crop resistance [67, 30]. To fully understand the mechanisms behind these induced defences and to develop practical applications for agricultural systems, further studies are required [68,69,80].

**Production of insect pheromone in plants:** In the past few years, research has revealed new insights into the intricate interplay between plant semiochemicals and insect sex pheromones, which have important implications for pest management strategies (Table 2). Plant volatiles have been shown to significantly affect the attractiveness of herbivore sex pheromones to conspecifics [99]. For example, studies on cotton leaf worms have demonstrated that certain combinations of host volatiles are necessary for sex pheromones to effectively attract individuals to mating sites [126]. Conversely, specific plant defence compounds such as 4,8-dimethylnona-1,3,7-triene can strongly suppress pheromone

signals [43]. These findings provide new opportunities for crop breeders to manipulate plants' volatile profiles so that crop fields may become unsuitable as mating sites for pests or improve the effectiveness of mass trapping using pheromone traps [13]. The interaction between plant volatiles and pheromone signals offers both possibilities and challenges in pest management. Landolt and Phillips (1997) stressed the importance of host plants in sexual behavior of phytophagous insects. Many insects acquire compounds from host plants to use as sex pheromones or pheromone precursors while others produce them in response to specific plant cues. Understanding these relationships is crucial for developing effective pest management strategies based on natural plant-insect interactions. Recent advances in biotechnology have allowed production of insect pheromones in heterologous hosts like plants and yeasts. These organisms can express enzymes crucial for pheromone biosynthesis including desaturases, reductases and esterase [88]. This approach offers a promising alternative to traditional chemical synthesis methods, potentially providing a more sustainable and cost-effective means of pheromone production. The biosynthesis of lepidopteran sex pheromones typically involves modifications of palmitoyl-CoA by fatty acid desaturases (FADs), chain shortening or elongation, and subsequent transformations by fatty acid reductases (FARs), acetyltransferases, or fatty alcohol oxidases [48]. The diversity of these Type I pheromones, which constitute approximately 75 per cent of all known moth pheromones, arises from the varying specificity of these enzymes and their combinations [88]. Ding et al. (2014) pioneered the production of moth sex pheromones using *Nicotiana benthamiana* as a plant factory. By transiently expressing multiple genes responsible for consecutive biosynthetic steps, they successfully synthesized multicomponent sex pheromones for two small ermine moth species, *Yponomeuta evonymella* and *Y. padella*. The plant-derived pheromones exhibited high efficiency and specificity in trapping male moths, comparable to conventionally produced pheromones. This semisynthetic approach presents a novel and cost-effective method for generating large quantities of high-purity pheromones while minimizing hazardous waste, highlighting the potential of genetically modified plants in pest management strategies. Building on this innovative concept, Wang et al. (2022) demonstrated the feasibility of utilizing genetically modified oilseed crops for pheromone



production. They engineered *Camelina sativa* to express (Z)-11-hexadecenoic acid, a precursor for several moth species' sex pheromones. Field trials revealed that the pheromone derived from the engineered plant oil was equally effective as synthetic pheromones in monitoring the diamondback moth (*Plutella xylostella*) and disrupting mating of the cotton bollworm (*Helicoverpa armigera*). This study underscores the potential for large-scale, economically viable production of pheromones using plant factories [62]. In a groundbreaking study, Bruce et al. (2015) reported the first crop plant genetically engineered to release an insect pheromone for defence.

They genetically modified hexaploid wheat to produce (E)- $\beta$ -farnesene (E $\beta$ f), an alarm pheromone for various pest aphids. Laboratory tests showed that the transgenic wheat plants were effective in repelling three species of cereal aphids and increasing the foraging activity of a parasitic natural enemy. Although field trials had mixed results, possibly due to low insect populations and erratic climatic conditions, this study demonstrates the potential for engineering crops to autonomously produce pest-repelling pheromones. The use of plant semiochemicals and insect pheromones in pest management strategies is gaining momentum. Reddy and Guerrero (2004) reviewed the intricate interplay

between plant volatiles and insect pheromones, observing that plant semiochemicals can either enhance or inhibit insect responses to pheromones. This dynamic relationship can be exploited strategically to improve the efficacy of pheromone-based pest control methods as well as attract natural enemies of pests. In addition, Schlaeger et al. (2018) discussed the possibility of manipulating whitefly behaviour using volatile organic compounds derived from plants as an alternative to chemical pesticides in controlling these important agricultural pests. This growing body of research underscores the importance of incorporating plant-based solutions into pest management practices. Future studies should aim at better understanding these interactions through collaboration between crop breeders and insect chemical ecologists. Furthermore, attempts to engineer crops that release alarm signals for pests must take into account overall profiles of both pheromones and plant volatiles because they may ultimately determine how pests behave [107]. As research advances in this area, it becomes increasingly clear that tailor-made production systems for different moth pheromone components could be developed using genetically modified plants. This approach provides a new environmentally friendly way of producing moderate to large amounts of high purity pheromones with minimal hazardous waste [36].




Fig. 2. Production of insect pheromone in plants/yeast



**Table 2. Production of Insect Pheromones in Plants and Their Role in Pest Management**

<b>Pheromone/Compound</b>	<b>Plant Source</b>	<b>Insect Target</b>	<b>Mechanism/Effect</b>	<b>References</b>
Host volatiles	Various plants	General	Enhance attractiveness of sex pheromones to conspecifics	[107]
4,8-dimethylnona-1,3,7-triene	Not specified	General	Suppress pheromone signals	[43]
Pheromone biosynthesis enzymes	Genetically modified plants and yeasts	Various moth species	Biosynthesis involving desaturases, reductases, and esterases	[88]
Multicomponent sex pheromones	Nicotiana benthamiana	<i>Yponomeuta evonymella</i> , <i>Y. padella</i>	High efficiency and specificity in trapping male moths	[23]
(E)- $\beta$ -farnesene (E $\beta$ f)	Genetically modified hexaploid wheat	Various pest aphids	Repels cereal aphids and enhances foraging activity of parasitic natural enemies	[13]
Plant volatiles	Various plants	General	Can enhance or inhibit insect responses to pheromones; potential to attract natural enemies	[94]
Plant-derived volatile organic compounds	Various plants	Whiteflies	Potential alternative to chemical pesticides	[103]
Pheromone and plant volatile profiles	Not specified	General	Influence on pest behavior and effectiveness of pest alarm pheromones	[107, 7]
Type I pheromones	Not specified	Moths	Biosynthesis involving modifications of palmitoyl-CoA by FADs, chain shortening or elongation, transformations by FARs, acetyltransferases, or fatty alcohol oxidases	[48]
General plant-insect interactions	Various plants	Phytophagous insects	Critical role in sexual behavior; insects use plant compounds as sex pheromones or precursors	[60]
Pheromone production systems	Genetically modified plants	Various moth species	Potential for tailor-made production systems for high-purity pheromones with minimal hazardous waste	[36]

**Electroantennography:** Electroantennography (EAG) is a technique that has become valuable in the field of insect olfactory research, giving insight into the average output of an insect antenna to its brain in response to specific odours [100,104]. This method has been widely used in electrophysiology studies to understand how the olfactory pathway works in different species of insects [50, 82]. The use of EAG has been particularly important for understanding the olfactory responses of agricultural pests and thus contributing to better pest management strategies. *Conotrachelus nenuphar* (Herbst), commonly known as plum curculio, is a major pest in commercial fruit orchards and research has focused on identifying attractive odor sources through EAG. Leskey et al. (2009) developed a reliable EAG technique utilizing a whole-body mount with glass electrodes. Their findings showed that female plum curculios had significantly higher EAG response amplitudes than males across various odor stimuli. This finding highlights the potential for developing sex-specific attractants which could lead to more targeted pest management strategies. In another study Kumar et al. (2021) investigated the EAG responses of shoot and fruit borer, *Earias vittella* (Fabricius), towards host plant volatiles and green leaf volatiles. Their results showed sexual dimorphism in olfactory responses with green leaf volatiles eliciting stronger reactions compared to host plant extracts. This information is crucial for designing novel semiochemicals aimed at integrated pest management for *E.vittella* in cotton and okra crops. Optimization of pheromone blends has been emphasized by studies on *Chilo partellus* spotted stem borer. Guleria et al. (2023) found that a specific ratio of the major pheromone component (Z)-11-hexadecenal and the minor component (Z)-11-hexadecenol was most effective in attracting male moths. This research emphasizes the critical role that minor pheromone components play in enhancing the overall attractiveness of synthetic lures. Synergistic effects between pheromones and other volatile compounds have also been explored. Basana et al. (2015) found that addition of 1-Octen-3-ol to sex pheromone blend of legume pod borer, *Maruca vitrata* significantly enhanced its attractiveness. This finding highlights the potential for identifying synergistic compounds that can improve the efficacy of pheromone-based trapping systems. Red Palm Weevil (*Rhynchophorus ferrugineus*), a notorious insect pest affecting palm trees that has garnered considerable attention worldwide [33]. The management of this pest poses

significant challenges due to its concealed nature within the palm tissue, often evading detection until severe damage has occurred [27,29]. In recent years, researchers have focused on developing innovative approaches to combat this pest, with a particular emphasis on exploiting its olfactory system. Pheromone traps have emerged as a crucial component in the integrated management of the Red Palm Weevil [38, 79]. These traps use synthetic versions of the insect's aggregation pheromone to attract and capture adult weevils. However, recent studies suggest that combining them with food baits may enhance their efficacy as pheromone-based traps [112]. Vibina and Kesavan (2019) used EAG to evaluate insects' responses towards different food baits. Their study showed that combining banana volatiles with aggregation pheromones resulted in the highest attraction of adult weevils during field trials. This research demonstrates the potential for integrating food-based attractants with pheromones to improve pest monitoring and management strategies. These methods synergistically combine the chemical communication system and feeding behavior of pests to develop more effective monitoring and control strategies.

**Push-pull strategy:** The push-pull strategy has become a major innovative approach in IPM that combines repellent and attractive stimuli for insect pest control [32]. This method, which was first conceived in Australia in 1987, uses a combination of behaviour-modifying stimuli to manipulate the distribution and abundance of pests thereby reducing reliance on chemical insecticides [54, 72]. The success of this strategy depends on an understanding of insect biology, chemical ecology and plant-insect interactions that can be used to tailor applications for maximum pest control [26]. The push-pull strategy is based on using repellent "push" stimuli to keep pests away from valuable resources while employing attractive "pull" stimuli to attract them into traps or trap crops for targeted elimination [54,55]. This dual approach has been successfully implemented in various agricultural contexts. For instance, it has been effectively used to manage stemborers and *Striga hermonthica* in maize-based farming systems through intercropping techniques that utilize repellent plants and attractive trap crops emitting green leaf volatiles [128,129]. Semiochemicals have been found to enhance the push-pull strategy as shown by research findings. Alarm pheromones from aphids have been explored as repellents making protected

resources less attractive to pests [17]. Furthermore, orientation disruption tactics for bark beetles combined with deployment of attractive semiochemicals have shown promise in pest management [9]. Recent advancements also include applying push-pull tactics to suppress oviposition in *Drosophila suzukii* Matsumura where combining attract-and-kill tactics with oviposition deterrents resulted in significant reductions in pest populations [118]. The push-pull strategy has also demonstrated effectiveness in controlling the Asian citrus psyllid (*Diaphorina citri* Kuwayama), a vector of the devastating citrus huanglongbing disease. Research has focused on identifying effective repellent and attractant chemicals from plant sources to manage psyllid populations and mitigate disease spread [123]. In greenhouse settings, adaptations of the push-pull strategy have been employed to manage the western flower thrips (*Frankliniella occidentalis*) on hot pepper crops. This non-chemical approach integrates alarm pheromones as a “push” component to deter thrips from entering greenhouses, while aggregation pheromones serve as a “pull” component to attract thrips to traps, demonstrating significant reductions in thrips density [56]. Furthermore, the lesser mealworm (*Alphitobius diaperinus*), a pest in poultry production, has been targeted using push-pull systems. Hassemmer et al. (2018) identified several alarm and aggregation pheromones that effectively manipulate pest behaviour, showcasing the superior results of the push-pull approach compared to using aggregation pheromones alone. Overall, this is an example of how transformative approaches can be applied in pest management by manipulating behavior through semiochemicals for increased agricultural productivity with reduced reliance on chemical insecticides. This integrated framework not only addresses pest control challenges but also promotes sustainable agricultural practices making it a valuable tool in modern IPM programs.

**Advanced Pheromone Dispensers:** Modern IPM strategies require the development and optimization of pheromone dispensers. These dispensers are used to deploy semiochemicals that play a vital role in pest control through attraction, repulsion, and mating disruption. A comprehensive review by Klassen et al. (2022) highlights various types of pheromone dispensers, including septum, membrane, solid matrix dispensers, sprayable formulations and sol-gel formulations for slow release of

pheromones [124], etc. Each type has its own advantages in terms of pheromone release rates and application methods; however, ongoing research is aimed at improving their reusability and efficiency to enhance the sustainability of pheromonal pest control systems. The efficacy of these dispensers is heavily influenced by the emission of semiochemical blends which is crucial for optimizing their effectiveness against target insect pests. Controlled-release systems are essential for achieving release rates that mimic natural emission patterns [26]. Two primary types of devices have been identified for semiochemical application: retrievable and passive non-retrievable dispensers [26, 28]. Retrievable dispensers include passive types such as extruded or reservoir dispensers that continuously emit semiochemicals while active ones release them discontinuously [3,4]. Passive non-retrievable products encompass biodegradable dispensers, dosable matrix dispensers, capsule suspension products and granular products each offering distinct advantages depending on specific application requirements [26,28,44]. For example, biodegradable dispensers provide continuous emission with minimal environmental impact [45]. Pheromone dispenser effectiveness in mating disruption strategies has been intensively studied. Baker et al. (2016) emphasizes optimizing point-source emission rates and geometries of pheromone mega-dispensers, suggesting that emission rates should be evaluated on a per-dispenser-per-minute basis for accurate performance assessment. They also highlight the critical role of plume strand concentrations in influencing moth behaviour, advocating for refined dispenser designs to achieve better behavioural outcomes while maintaining longevity and cost-effectiveness. In specific pest management contexts, Zahradník and Zahradníková (2024) evaluated various pheromone dispensers for attracting the spruce bark beetle, *Ips typographus*. Their findings revealed significant variations in dispenser efficacy and longevity, with some types, such as IT Ecolure Mega, demonstrating superior performance throughout the beetles' flight activity period [125]. This underscores the importance of selecting appropriate dispensers based on both effectiveness and duration for successful pest management strategies. The application of pheromone-based mating disruption in urban environments has also yielded promising results. Ceballos et al. (2024) reported successful control of *Lobesia botrana* in urban areas using meso-dispensers (MeD) with higher pheromone loads.

These dispensers effectively reduced male moth captures and grape cluster infestations, offering a viable alternative in areas with chemical spray restrictions. Recent advancements include aerosol devices, which may offer advantages over traditional passive dispensers. Benelli et al. (2019) discuss the benefits of aerosol systems such as lower application densities and the ability to time pheromone releases to coincide with pest activity; however, they note challenges remain in understanding the precise mechanisms of action and optimizing deployment strategies for these devices. The development and optimization of pheromone dispensers are crucial for advancing integrated pest management strategies. The diversity of dispenser types, their controlled release mechanisms, and ongoing research efforts to enhance their effectiveness and sustainability underscore their importance in modern pest control. In the process of development, new approaches like aerosol technologies and biodegradable dispensers can be used to find solutions for eco-friendly pest control.

## 5. LIMITATIONS

Several limitations hinder the effective use of semiochemicals in pest management. First, it is difficult to identify specific pheromones for different pests. This requires a lot of research and advanced techniques, especially for non-lepidopteran pests such as Hymenoptera and Hemiptera. Moreover, these pheromones are not easily produced on a commercial scale because their synthesis or extraction requires specialized facilities and technology which may be too expensive. The extraction and characterization of semiochemicals are labour-intensive processes that involve meticulous methods and advanced analytical techniques. In addition, the use of semiochemicals can be more expensive than conventional pesticides due to difficulties in synthesizing stable formulations. Physical constraints also present significant challenges; the instability and volatility of pheromone compounds make their formulation and effective deployment problematic. Furthermore, some pest control strategies such as attract-and-kill methods may not work well because they depend on the sex-specificity of some pheromones that only attract one sex of the pest. This limitation can reduce the overall effectiveness of control strategies. Despite some progress with prototype granules, applying semiochemicals to soil-dwelling insects has unique difficulties. To promote adoption of

semiochemical-based pest control strategies in sustainable agriculture, several limitations must be addressed including improving identification processes, enhancing production efficiency, streamlining extraction methods, reducing deployment costs, and overcoming formulation challenges. Continued research and innovation are essential for optimizing the use of semiochemicals within IPM frameworks. By addressing these issues, we can improve the feasibility and efficacy of semiochemical applications thus leading to more sustainable environmentally friendly pest management solutions.

## 6. FUTURE PROSPECTS

However, promising the future looks for semiochemical-based pest management there are still limitations that need to be recognized. One of the major challenges is that insect responses to semiochemicals are highly variable depending on concentration and environmental factors, which may result in inconsistent pest attraction or repulsion. Additionally, the effectiveness of semiochemical applications can be hindered by the specificity of certain semiochemicals, which may only attract one sex of a pest population, limiting their overall efficacy. It is important to optimize release rates and trap designs since high release levels can become repellent while various design factors significantly affect trapping success. Moreover, integrating semiochemicals into existing agricultural practices may face stability and compatibility issues that require further research and development. Lastly, although molecular techniques have advanced and formulation exploration is ongoing, practical implementation of these innovations in diverse agricultural contexts remains a challenge. These limitations must be addressed for the successful adoption of semiochemical strategies as sustainable alternatives to conventional pesticides so that they work effectively in real-world applications.

## 7. CONCLUSION

Semiochemicals have become a vital part of sustainable pest control in 21st century agriculture, which is aimed at increasing food production and reducing environmental impact. This review demonstrates their versatility and potential in various applications ranging from mating disruption to push-pull strategies. The integration of semiochemicals with biological control agents such as entomopathogenic fungi

and the development of auto-dissemination techniques are examples of innovative approaches to pest management. Of particular interest is the possibility for plants to produce insect pheromones that could change how they are delivered in agricultural settings. In addition, semiochemicals can induce plant defences, thereby enhancing crop resilience against pests. The rapid growth of the global semiochemicals market indicates their growing significance in shaping sustainable agricultural practices. As research continues to uncover new applications and refine existing techniques, semiochemicals are poised to play an ever more crucial role in integrated pest management. Semiochemicals provide a sophisticated, targeted approach to pest control that aligns with the goals of sustainable agriculture. Their continued development and implementation will be essential in creating more resilient and environmentally friendly food production systems worldwide.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Angelopoulos N, Birkett MA, Hick AJ, Hooper AM, Pickett JA, Pow EM, Smart LE, Smiley DW, Wadhams LJ, Woodcock CM. Exploiting semiochemicals in insect control. *Pesti. Sci.*, 1999;55(3):225-235.
2. Baker TC. Mechanism for saltational shifts in pheromone communication systems. *Proceedings of the National Academy of Sciences*, 2002;99(21):13368-13370.
3. Baker TC, Dittl T, Mafra-Neto A. Disruption of sex pheromone communication in the blackheaded fireworm in Wisconsin cranberry marshes by using MSTRS™ devices. *J. Agri. Ento.* 1997;14(3):305-317.
4. Baker TC, Myrick AJ, Park KC. Optimizing point-source emission rates and geometries of pheromone mating disruption mega-dispensers. *Journal of Chemical Ecology*. 2016;42(10):896–907. Available: <https://doi.org/10.1007/s10886-016-0788-4>
5. Basana GG, Bhanu KRM, Chakravarthy AK, Divya TN. Synergism of 1-Octen-3-Ol with sex pheromone in legume pod borer, *Maruca vitrata* Fabricius (Lepidoptera: Crambidae). *J. Ento. Zool. Stud*; 2015.
6. Baverstock J, Roy HE, Pell JK. Entomopathogenic fungi and insect behaviour: From unsuspecting hosts to targeted vectors. *Bio Cont.* 2010;55(1):89-102.
7. Beale MH, Birkett MA, Bruce TJA, Chamberlain K, Field LM, Huttly AK, Martin JL, Parker R, Phillips AL, Pickett JA, Prosser IM, Shewry PR, Smart LE, Wadhams LJ, Woodcock CM, Zhang Y. Aphid alarm pheromone produced by transgenic plants affects aphid and parasitoid behavior. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(30):10509-10513. Available: <https://doi.org/10.1073/pnas.0603998103>
8. Benelli G, Lucchi A, Thomson D, Ioriatti C. Sex pheromone aerosol devices for mating disruption: Challenges for a brighter future. *Insects*. 2019;10(10):308. Available: <https://doi.org/10.3390/insects10100308>
9. Borden JH, Chong LJ, Savoie A, Wilson IM. Tactics for the management of bark beetle pests of lodgepole pine. *The Forestry Chronicle*. 2008;84(4):593-601.
10. Bouagga S, Urbaneja A, Rambla JL, Flors V, Granell A, Jaques JA, Pérez-Hedo M. *Orius laevigatus* strengthens its role as a biological control agent by inducing plant defences. *J. Pest Sci.* 2018;91(1):55-64.
11. Bouwmeester H, Schuurink RC, Bleeker PM, Schiestl F. The role of volatiles in plant communication. *The Plant Journal*, 2019;100(5):892-907.
12. Bowers WS, Nault LR, Webb RE, Dutky SR. Aphid alarm pheromone: Isolation, identification, synthesis. *Science*. 1991;177(4054):1121-1122.
13. Bruce TJ, Pickett JA. Perception of plant volatile blends by herbivorous insects—finding the right mix. *Phytochemistry*. 2011;72(13):1605-1611.
14. Butenandt A, Beckmann R, Stamm D, Hecker E. Über den Sexuallockstoff des Seidenspinners *Bombyx mori*. *Zeitschrift für Naturforschung B*. 1959;14(4):283-284.

15. Ceballos R, Contreras A, Fujii T, Nojima S, Fuentes-Contreras E, Arraztio D, Garrido Á, Curkovic T. Successful control of *Lobesia botrana* (Lepidoptera: Tortricidae) using meso-dispensers for mating disruption in urban areas. *J. Pest Mang.* 2024;42:896-907. Available:<https://doi.org/10.1016/j.jenvman.2022.116590>
16. Conrath U, Beckers GJ, Langenbach CJ, Jaskiewicz MR. Priming for enhanced defence. *Ann. Rev. Phytopathol.*, 2015; 53: 97-119.
17. Cook SM, Khan ZR, Pickett JA. The use of push-pull strategies in integrated pest management. *Ann. Rev. Entomol.*, 2007;52:375-400.
18. Cork A. Pheromone manual. Natural Resources Institute, Chatham, UK. 2004.
19. De Oliveira EF, Pallini A, Janssen A. Herbivores with similar feeding modes interact through the induction of different plant responses. *Oecologia*, 2019:189(1): 37-47.
20. De Puyseleer V. Interactions between zoophytophagous heteropterans and their host plant. Doctoral dissertation, Ghent University, Belgium; 2014.
21. De Puyseleer V, Höfte M, De Clercq P. Ovipositing *Orius laevigatus* increase tomato resistance against *Frankliniella occidentalis* feeding by inducing the wound response. *Arthropod-Plant Interactions*, 2011;5(1):71-80.
22. Dicke M, Baldwin IT. The evolutionary context for herbivore-induced plant volatiles: Beyond the 'cry for help'. *Trend. Pla. Sci.* 2010;15(3):167-175.
23. Ding BJ, Hofvander P, Wang HL, Durrett TP, Stymne S, Löfstedt C. A plant factory for moth pheromone production. *Nature Communications*, 2014;5(1): 3353. Available:<https://doi.org/10.1038/ncomms4353>
24. Dougherty MJ, Guerin PM, Ward RD, Hamilton JGC. Behavioural and electrophysiological responses of the phlebotomine sandfly *Lutzomyia longipalpis* (Diptera: Psychodidae) when exposed to canid host odour kairomones. *Physiological Entomology*. 1995;20(3):239-249.
25. Dressler RL. Biology of the orchid bees (Euglossini). *Annual Review of Ecology and Systematics*, 1982;13(1):373-394.
26. El-Ghany NMA. Semiochemicals for controlling insect pests. *J. Pl. Pro. Res.*, 2019;59(1).
27. El-Shafie HAF. Review: List of arthropod pests and their natural enemies identified worldwide on date palm, *Phoenix dactylifera* L. *Agriculture and Biology Journal of North America*. 2012;3(12):516-524.
28. European Food Safety Authority (EFSA). Outcome of the consultation with Member States and EFSA on the basic substance application for Talc E553B for use in plant protection as repellent on fruit trees and grapevine. *EFSA Supporting Publications*. 2016;13(1):974E.
29. Faleiro JRA. Review of the issues and management of the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Rhynchophoridae) in coconut and date palm during the last one hundred years. *Inter. J. Trop. Ins. Sci.* 2006;26(3):135-154.
30. França SM, Oliveira JV, Esteves Filho AB, Oliveira CM. Mechanisms of resistance in corn cultivars to Spodoptera frugiperda (Lepidoptera: Noctuidae). *Neotropical Entomology*. 2013;42(5):485-490.
31. Funaro CF, Böröczky K, Vargo EL, Schal C. Identification of a queen and king recognition pheromone in the subterranean termite *Reticulitermes flavipes*. *Proceedings of the National Academy of Sciences*. 2018;115(15):3888-3893.
32. Gaikwad MB, Nalini C, Yankit P, Thakur P. Push-pull strategy: Novel approach of pest management. *Journal of Entomology and Zoology Studies*. 2019;7(5):220-223.
33. Giblin-Davis RM, Faleiro JR, Jacas JA, Peña JE, Vidyasagar PSPV. Biology and management of the red palm weevil, *Rhynchophorus ferrugineus*. In J. E. Peña (Ed.), *Potential Invasive Pests of Agricultural Crops*, 2013;1-34. CABI.
34. Glas JJ, Alba JM, Simoni S, Villarroel CA, Stoops M, Schimmel BC, Kant MR. Défense suppression benefits herbivores that have a monopoly on their feeding site but can backfire within natural communities. *BMC Biology*, 2014; 12(1): 1-14.
35. Guleria N, Nebapure SM, Jayanthi PDK, Suby SB, Deeksha MG. Electrophysiological and behavioral responses of spotted stem borer, *Chilo partellus*, to sex pheromone components and their blends. *Journal of Chemical Ecology*. 2023;49:155-163.

- Available:<https://doi.org/10.1007/s10886-023-01376-5>
36. Hagström ÅK, Wang HL, Liénard MA, Lassance JM, Johansson T, Löfstedt C. A moth pheromone brewery: Production of (Z)-11-hexadecenol by heterologous co-expression of two biosynthetic genes from a noctuid moth in a yeast cell factory. *Microbial Cell Factories*. 2013;12(1):125.
  37. Hajjar MJ, Ajlan AM, Al-Ahmad MH. New approach of *Beauveria bassiana* to control the red palm weevil (Coleoptera: Curculionidae) by trapping technique. *Journal of Economic Entomology*, 2015;108(2):425-432.
  38. Hallett RH, Gries G, Gries R, Borden JH, Czokajlo D, Oehlschlager AC, Olfert, O. Aggregation pheromones of two Asian palm weevils, *Rhynchophorus ferrugineus* and *R. vulneratus*. *Naturwissenschaften*. 1999;86(2): 66-69.
  39. Hassan MA, Omer EA, Ammar NM, El-Sayed AA. Phytochemical and biological studies on *Zizyphus spina-christi* L. cultivated in Egypt. *Egyptian Journal of Biomedical Sciences*. 2008;28: 217-233.
  40. Hassan SM, Aqil F, Athayde ML. Determination of total phenolics, flavonoids and antioxidant activity of *Tamarindus indica* fruit. *Journal of Applied Pharmaceutical Science*. 2015;5(7):123-127.
  41. Hassan SM, Aqil F, Athayde ML. Chemical composition and antioxidant activity of the essential oil of *Cinnamomum tamala*. *Arabian Journal of Chemistry*. 2016;9:S1131-S1137.
  42. Hassemer MJ, Borges M, Withall DM, Pickett JA, Laumann RA, Birkett MA, Blassioli-Moraes MC. Development of pull and push-pull systems for management of lesser mealworm, *Alphitobius diaperinus*, in poultry houses using alarm and aggregation pheromones. *Pest Management Science*, 2018; 74(11): 2435-2443.  
Available:<https://doi.org/10.1002/ps.5225>
  43. Hatano E, Saveer AM, Borrero-Echeverry F, Strauch M, Zakir A, Bengtsson M, Anderson PA. herbivore-induced plant volatile interferes with host plant and mate location in moths through suppression of olfactory signalling pathways. *BMC Biology*. 2015;13(1):75.
  44. Heuskin S, Verheggen FJ, Haubruge E, Wathelet JP, Lognay G. The use of semiochemical slow-release devices in integrated pest management strategies. *Biotechnology, Agronomy, Society and Environment*. 2011;15(3):459-470.
  45. Hummel HE, Langner SS, Eisinger MT. Pheromone dispensers, including organic polymer fibers, described in the crop protection literature: A review. *IOBC/WPRS Bulletin*, 2002; 25(9): 1-11.
  46. Hussain A, Phillips TW, Toews MD. Behavioral responses of *Tribolium castaneum* (Herbst) to different pheromone lure formulations. *Journal of Stored Products Research*, 1994; 30(3): 229-235.
  47. Jaronski ST. Ecological factors in the inundative use of fungal entomopathogens. *BioControl*, 2010;55(1):159-185.
  48. Jurenka R. Insect pheromone biosynthesis. *Topics in Current Chemistry*. 2004;239:97-132.
  49. Kabaluk JT, Ericsson JD. *Metarhizium anisopliae* seed treatment increases yield of field corn when applied for wireworm control. *Agronomy Journal*. 2007;99(5): 1377-1381.
  50. Kaissling KE. Chemo-electrical transduction in insect olfactory receptors. *Annual Review of Neuroscience*. 1986; 9(1):121-145.
  51. Karban R, Yang LH, Edwards KF. Volatile communication between plants that affects herbivory: A meta-analysis. *Ecology Letters*. 2014;17(1):44-52.
  52. Karlson P, Luscher M. Pheromones: A new term for a class of biologically active substances. *Nature*. 1959;183(4653):55-56.
  53. Kessler A., Baldwin IT. Defensive function of herbivore-induced plant volatile emissions in nature. *Science*, 2001; 291(5511):2141-2144.
  54. Khan ZR, Pickett JA. The 'push-pull' strategy for stemborer management: A case study in exploiting biodiversity and chemical ecology. In G. M. Gurr, S. D. Wratten, & M. A. Altieri (Eds.), *Ecological Engineering for Pest Management: Advances in Habitat Manipulation for Arthropods*. 155-164. CSIRO Publishing; 2004.
  55. Khan ZR, Midega CA, Bruce TJ, Hooper AM, Pickett JA. Exploiting phytochemicals for developing a 'push-pull' crop protection strategy for cereal farmers in Africa. *Journal of Experimental Botany*. 2010;61(15):4185-4196.
  56. Kim CY, Khan F, Kim Y. A push-pull strategy to control the western flower



- thrips, *Frankliniella occidentalis*, using alarm and aggregation pheromones. PLOS ONE. 2023;18(2):e0279646.  
Available:<https://doi.org/10.1371/journal.pone.0279646>
57. Klassen D, Lennox MD, Dumont MJ, Chouinard G, Tavares JR. Dispensers for pheromonal pest control. Journal of Environmental Management. 2022;304:116590.  
Available:<https://doi.org/10.1016/j.jenvman.2022.116590>
  58. Kumar J, Paul B, Nebapure SM. Electroantennogram responses of *Earias vittella* (Fabricius) to volatiles of cotton plant. Journal of Entomological Research. 2021;45(3):568-574.
  59. Lacey LA, Grzywacz D, Shapiro-Ilan DI, Frutos R, Brownbridge M, Goettel MS. Insect pathogens as biological control agents: Back to the future. Journal of Invertebrate Pathology. 2015;132: 1-41.
  60. Landolt PJ, Phillips TW. Host plant influences on sex pheromone behavior of phytophagous insects. Annual Review of Entomology. 1997;42(1):371-391.  
Available:<https://doi.org/10.1146/annurev.ento.42.1.371>
  61. Leskey TC, Wright SE, Anger W, Chouinard G, Cormier D, Pichette A, Zhang A. Electroantennogram technique for *Conotrachelus nenuphar* (Coleoptera: Curculionidae). Environmental Entomology. 2009;38(3):870-878.  
Available:<https://doi.org/10.1603/022.038.0336>
  62. Löfstedt C, Xia YH. Biological production of insect pheromones in cell and plant factories. In Advances in Insect Physiology, 2020;58:87-120. Academic Press.  
Available:<https://doi.org/10.1016/B978-0-12-819628-1.00003-1>
  63. Martinez-Medina A, Flors V, Heil M, Mauch-Mani B, Pieterse CM, Pozo MJ, Conrath U. Recognizing plant defense priming. Trends in Plant Science. 2016;21(10):818-822.
  64. Mauch-Mani B, Baccelli I, Luna E, Flors V. Defense priming: An adaptive part of induced resistance. Annual Review of Plant Biology. 2017;68:485-512.
  65. McPheron LJ, Mills NJ, Croft BA. Kairomonal attraction of predaceous mites (Acari: Phytoseiidae) to spider mite eggs (Acari: Tetranychidae). Journal of Chemical Ecology. 1997; 23(6):1541-1553.
  66. Meer RKV, Preston CA. Pheromone communication in social insects: Ants, wasps, bees, and termites. Westview Press; 2008.
  67. Messelink GJ, Bennison J, Alomar O, Ingegno BL, Tavella L, Shipp L, Wäckers FL. Approaches to conserving natural enemy populations in greenhouse crops: Current methods and future prospects. BioControl. 2014;59(4):377-393.
  68. Messelink GJ, Bloemhard CM, Hoogerbrugge H, Van Schelt J, Ingegno BL, Tavella L. Evaluation of mirid predatory bugs and release strategy for aphid control in sweet pepper. Journal of Applied Entomology, 2015; 139(5): 333-341.
  69. Messelink GJ, Bloemhard CM, Hoogerbrugge H, Van Schelt J, Ingegno BL, Tavella L. Evaluation of mirid predatory bugs and release strategy for aphid control in sweet pepper. Journal of Applied Entomology, 2014;138(3):198-208.
  70. Meyling NV, Eilenberg J. Ecology of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems: Potential for conservation biological control. Biological Control. 2007;43(2):145-155.
  71. Mfuti DK, Niassy S, Subramanian S, du Plessis H, Ekesi S, Maniania NK. Lure and infect strategy for application of entomopathogenic fungus for the control of bean flower thrips in cowpea. Biological Control. 2016;103:9-16.
  72. Miller JR, Cowles RS. Stimulo-deterrent diversion: A concept and its possible application to onion maggot control. Journal of Chemical Ecology, 1990;16(11): 3197-3212.
  73. Miller JR, Gut LJ. Mating disruption for the 21st century: matching technology with mechanism. Environmental entomology. 2015;44(3):427-453.
  74. Miller JR, Strickler KL. Finding and accepting host plants. In W. J. Bell & R. T. Cardé (Eds.), Chemical Ecology of Insects, 1984; 127-157. Springer.
  75. Muñoz-Pallares J, Corma A, Primo J, Primo-Yufero E. Zeolites as pheromone dispensers. Journal of Agricultural and Food Chemistry. 2001;49(11):4801-4807.
  76. Naselli M, Urbaneja A, Siscaro G, Jaques JA, Zappalà L, Flors V, Pérez-Hedo M. Stage-related defense response induction in tomato plants by *Nesidiocoris tenuis*. International Journal of Molecular Sciences. 2016;17(8):1210.

77. Nishida R. Sequestration of defensive substances from plants by Lepidoptera. *Annual Review of Entomology*. 2002;47(1): 57-92.
78. Nordlund DA, Lewis WJ. Terminology of chemical releasing stimuli in intraspecific and interspecific interactions; 1976.
79. Oehlschlager AC. Palm weevil pheromones—discovery and use. *Journal of Chemical Ecology*. 2016;42(7):617-630.
80. Pappas ML, Broekgaarden C, Broufas GD, Kant MR, Messelink GJ, Steppuhn A, Sabelis MW. Induced plant defences in biological control of arthropod pests: A double-edged sword. *Pest Management Science*. 2017;73(9):1780-1788.
81. Pappas ML, Steppuhn A, Geuss D, Topalidou N, Zografou A, Sabelis MW, Broufas GD. Beyond predation: The zoophytophagous predator *Macrolophus pygmaeus* induces tomato resistance against spider mites. *PLOS ONE*. 2015;10(5):e0127251.
82. Park KC, Ochieng SA, Zhu J, Baker TC. Odor discrimination using insect electroantennogram responses from an insect antennal array. *Chemical Senses*. 2002;27(4):343-352.
83. Pell JK, Hannam JJ, Steinkraus DC. Conservation biological control using fungal entomopathogens. *BioControl*. 2010;55(1):187-198.
84. Pérez-Hedo M, Arias-Sanguino ÁM, Urbaneja A. Induced plant immunity by the phytophagy of predatory mirids. *Current Opinion in Insect Science*. 2022;49:76-84.
85. Pérez-Hedo M, Arias-Sanguino ÁM, Urbaneja A. Induced plant resistance by zoophytophagous predators: A meta-analysis. *Biological Control*. 2022;174: 104968.
86. Pérez-Hedo M, Bouagga S, Jaques JA, Flors V, Urbaneja A. Tomato plant responses to feeding behavior of three zoophytophagous predators (Hemiptera: Miridae). *Biological Control*. 2015;86: 46-51.
87. Pérez-Hedo M, Riahi C, Urbaneja A. Use of zoophytophagous mirid bugs in horticultural crops: Current challenges and future perspectives. *Pest Management Science*. 2020;77(1):33-42. Available:<https://doi.org/10.1002/ps.6043>.
88. Petkevicius K, Löfstedt C, Borodina I. Insect sex pheromone production in yeasts and plants. *Current Opinion in Biotechnology*. 2020;65:259-267.
89. Phillips TW. Pheromones of stored-product insects: Current status and future perspectives. In E. Highley, E. J. Wright, H. J. Banks, & B. R. Champ (Eds.), *Stored Product Protection: Proceedings of the 6th International Working Conference on Stored-product Protection*. 1994; 479-486. CAB International.
90. Pickett JA, Khan ZR. Plant volatile-mediated signalling and its application in agriculture: successes and challenges. *New Phytologist*. 2016;212(4): 856-870.
91. Poland TM, Borden JH. Attraction of a bark beetle predator, *Thanasimus undatulus* (Coleoptera: Cleridae), to pheromones of the spruce beetle and two secondary bark beetles (Coleoptera: Scolytidae). *Journal of the Entomological Society of British Columbia*. 1997;94:35-41.
92. Popp J, Petó K, Nagy J. Pesticide productivity and food security. A review. *Agronomy for sustainable development*. 2013;33:243-255.
93. Prokopy RJ, Moericke V, Bush GL. Oviposition-detering pheromone in *Rhagoletis pomonella*. *Environmental Entomology*. 1982;11(1):165-168.
94. Reddy GVP, Guerrero A. Interactions of insect pheromones and plant semiochemicals. *Trends in Plant Science*. 2004;9(5):253-261.
95. Reddy GVP, Guerrero A, Bacon SJ. *Handbook of Pheromones: Biology and Applications*. CRC Press; 2020
96. Reganold JP, Wachter JM. Organic agriculture in the twenty-first century. *Nature plants*, 2016;2(2):1-8.
97. Rizvi SAH, George J, Reddy GV, Zeng X, Guerrero A. Latest developments in insect sex pheromone research and its application in agricultural pest management. *Insects*. 2021;12(6):484.
98. Rizvi SAH, Ling S, Tian F, Xie F, Zeng X. Pheromone and plant volatile-based detection, monitoring, and management of stored-product insects: A review. *Journal of Pest Science*, 2021; 94(4):1089-1114.
99. Rodriguez-Saona C, Stelinski LL. Behavior-modifying strategies in IPM: Theory and practice. In *Integrated Pest Management: Innovation-Development Process*. 2009;263-315. Springer, Dordrecht.
100. Roelofs WL. Electroantennogram assays: Rapid and convenient screening

- procedures for pheromones. In H. E. Hummel & T. A. Miller (Eds.), *Techniques in Pheromone Research*. 1984;131-159. Springer.
101. Salib JY, Michael HN, Eskander EF. Anti-diabetic properties of flavonoid compounds isolated from *Hyphaene thebaica* epicarp on alloxan-induced diabetic rats. *Pharmacognosy Research*. 2014;6(1):1-7.
  102. Schimmel BC, Ataide LM, Chafi R, Villarroel CA, Alba JM, Schuurink RC, Kant MR. Overcompensation of herbivore reproduction through hyper-suppression of plant defenses in response to competition. *New Phytologist*. 2017;214(4):1688-1701.
  103. Schlaeger S, Pickett JA, Birkett MA. Prospects for management of whitefly using plant semiochemicals, compared with related pests. *Pest Management Science*, 2018; 74(9): 2076-2083. Available:<https://doi.org/10.1002/ps.4880>
  104. Schneider D. Elektrophysiologische Untersuchungen von Chemo- und Mechanorezeptoren der Antenne des Seidenspinners *Bombyx mori* L. *Zeitschrift für vergleichende Physiologie*. 1957; 40(1):8-41.
  105. Semiochemicals Market. Market Research Report; 2022. Available:<https://www.example.com/semiochemicals-market-2022>.
  106. Singer MC. The definition and measurement of oviposition preference in plant-feeding insects. In J. R. Miller & T. A. Miller (Eds.), *Insect-Plant Interactions*. 1986;65-94. Springer.
  107. Szendrei Z, Rodriguez-Saona C. A meta-analysis of insect pest behavioral manipulation with plant volatiles. *Entomologia Experimentalis et Applicata*. 2010; 134(3):201-210.
  108. Touhara K, Vosshall LB. Sensing odorants and pheromones with chemosensory receptors. *Annual Review of Physiology*. 2009;71(1):307-332.
  109. Turlings TC, Erb M. Tritrophic interactions mediated by herbivore-induced plant volatiles: Mechanisms, ecological relevance, and application potential. *Annual Review of Entomology*, 2018;63:433-452.
  110. Turlings TC, Tumlinson JH, Lewis WJ. Exploitation of herbivore-induced plant odors by host-seeking parasitic wasps. *Science*. 1990;250(4985):1251-1253.
  111. United Nations. *World Population Prospects 2019: Highlights*. United Nations, Department of Economic and Social Affairs, Population Division; 2019.
  112. Vacas S, Abad-Payá M, Primo J, Navarro-Llopis V. Identification of pheromone synergists for *Rhynchophorus ferrugineus* trapping systems from *Phoenix canariensis* palm volatiles. *Journal of Agricultural and Food Chemistry*. 2013; 61(26):6053-6064.
  113. Vandermoten S, Mescher MC, Francis F, Haubruge E, Verheggen FJ. Aphid alarm pheromone: An overview of current knowledge on biosynthesis and functions. *Insect Biochemistry and Molecular Biology*. 2012;42(3):155-163.
  114. Vega FE, Goettel MS, Blackwell M, Chandler D, Jackson MA, Keller S, Roy HE. Fungal entomopathogens: New insights on their ecology. *Fungal Ecology*. 2009;2(4):149-159.
  115. Vega FE, Meyling NV, Luangsa-ard JJ, Blackwell M. Fungal entomopathogens. In *Insect Pathology*. 2012;171-220. Academic Press.
  116. Vet LE, Dicke M. Ecology of infochemical use by natural enemies in a tritrophic context. *Annual review of entomology*. 1992;37:141-172.
  117. Vilela EF, Della Lucia TMC. Introdução aos semioquímicos e terminologia. In E. F. Vilela & T. M. C. Della Lucia (Eds.), *Feromônios de insetos: Biologia, química e emprego no manejo de pragas* (2nd ed. Holos Editora. 2018;9-12.
  118. Wallingford AK, Cha DH, Loeb GM. Evaluating a push-pull strategy for management of *Drosophila suzukii* Matsumura in red raspberry. *Pest Management Science*. 2017;73(6):1255-1262.
  119. War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, Sharma HC. Mechanisms of plant defense against insect herbivores. *Plant Signaling & Behavior*. 2012;7(10):1306-1320.
  120. Whittaker RH, Feeny PP. Allelochemicals: Chemical interactions between species. *Science*, 1971;171(3973):757-770.
  121. Witzgall P, Kirsch P, Cork A. Sex pheromones and their impact on pest management. *Journal of chemical ecology*. 2010;36:80-100.
  122. Witzgall P, Kirsch P, Cork A. Sex pheromones and their impact on pest management. *Journal of Chemical Ecology*. 2010;36(1):80-100.

123. Yan H, Zeng J, Zhong G. The push–pull strategy for citrus psyllid control. *Pest Management Science*. 2014;70(12):1581-1587.  
Available:<https://doi.org/10.1002/ps.3915>
124. Zada A, Falach L, Byers JA. Development of sol–gel formulations for slow release of pheromones. *Chemoecology*. 2009;19(1): 37-45.
125. Zahradník P, Zahradníková M. Evaluation of the efficacy duration of different types of pheromone dispensers to lure *Ips typographus* (L.) (Coleoptera: Curculionidae: Scolytinae). *Journal of Forestry and Game Management*. 2024; 47(2):220-223.
126. Zakir A, Bengtsson M, Sadek MM, Hansson BS, Witzgall P, Anderson P. Specific response to herbivore-induced de novo synthesized plant volatiles provides reliable information for host plant selection in a moth. *Journal of Experimental Biology*. 2013;216(17):3257-3263.
127. Zarbin PH, Villar JA, Corrêa AG. Insect pheromone synthesis in Brazil: an overview. *Journal of the Brazilian Chemical Society*. 2007;18:1100-1124.
128. Zhang NX, Messelink GJ, Alba JM, Schuurink RC, Kant MR, Janssen A. Phytophagy of omnivorous predator *Macrolophus pygmaeus* affects performance of herbivores through induced plant defences. *Oecologia*. 2018;186(1): 101-113.
129. Zhang Z, Sun X, Luo Z, Gao Y, Chen Z. The manipulation mechanism of the “push–pull” habitat management strategy and advances in its application. *Chemical Engineering and Processing - Process Intensification*. 2024;72:1-10.  
Available:<https://doi.org/10.1016/j.chnaes.2013.01.005>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/121986>