



The Large Scale Universe as a Quasi Quantum White Hole

U. V. S. Seshavatharam^{1*}, Eugene Terry Tatum² and S. Lakshminarayana³

¹Honorary Faculty, I-SERVE, Survey no-42, Hitech city, Hyderabad-84, Telangana, India.

²760 Campbell Ln. Ste 106 #161, Bowling Green, KY, USA.

³Department of Nuclear Physics, Andhra University, Visakhapatnam-03, AP, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author UVSS designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors ETT and SL managed the analyses of the study. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. David Garrison, University of Houston-Clear Lake, USA.

(2) Professor. Hadia Hassan Selim, National Research Institute of Astronomy and Geophysics, Egypt.

Reviewers:

(1) Abhishek Kumar Singh, Magadh University, India.

(2) Mohsen Lutephy, Azad Islamic university (IAU), Iran.

(3) Sie Long Kek, Universiti Tun Hussein Onn Malaysia, Malaysia.

(4) N.V.Krishna Prasad, GITAM University, India.

(5) Maryam Roushan, University of Mazandaran, Iran.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/66092>

Original Research Article

Received 17 January 2021

Accepted 23 March 2021

Published 01 April 2021

ABSTRACT

We emphasize the point that, standard model of cosmology is basically a model of classical general relativity and it seems inevitable to have a revision with reference to quantum model of cosmology. Utmost important point to be noted is that, 'Spin' is a basic property of quantum mechanics and 'rotation' is a very common experience. In this context, we propose five assumptions in line with Planck mass as the baby universe. We appeal that, 1) Universe can be modelled as a time-reversed black hole (a white hole) with rotation and light speed expansion, and 2) 'Light speed expanding cosmic space' can be called as 'Flat space'. With reference to light speed expansion, if one is willing to re-define cosmic red shift as $[z/(1+z)]$, without considering Lambda cosmology model of matter density fractions, light travel distances can be reproduced with marginal error. Advantages of our assumptions are, 1) A quantum model of cosmology can be developed with unification of general

*Corresponding author: E-mail: seshavatharam.uvs@gmail.com;

theory of relativity and quantum mechanics. 2) Tension in estimating the current Hubble parameter can be eliminated via scaled Hawking's black hole temperature formula with great confidence. 3) Galactic dark matter and visible matter can be studied in a unified manner. 4) Galactic light travel distances can be estimated very easily without matter density fractions. 5) Big bang and inflation like non-general relativistic concepts can be relinquished with further study.

Keywords: white hole cosmology; Planck mass as the baby universe; cosmic rotation; light speed expansion; dark matter; visible matter; galactic radii; galactic angular velocity.

LIST OF SYMBOLS

At any stage of cosmic evolution,

- 1) Cosmic time = t
- 2) Cosmic Hubble parameter = H_t
- 3) Cosmic angular velocity = ω_t
- 4) Ratio of Hubble parameter to angular velocity = Y_t
- 5) Cosmic radius = R_t
- 6) Cosmic total mass = M_t
- 7) Cosmic temperature = T_t
- 8) Galactic star flat rotation speed = V_{Gt}
- 9) Galactic angular velocity = $(\omega_G)_t$
- 10) Visible mass of galaxy = $(M_{Gv})_t$
- 11) Dark mass of galaxy = $(M_{Gd})_t$
- 12) Total mass of galaxy = $(M_G)_t$
- 13) Time dependent dark-visible reference mass unit = M_{Xt}
- 14) Radius of galaxy = $(R_G)_t$
- 15) Angular velocity of galaxy = $(\omega_G)_t$
- 16) Wavelength of light received from observed galaxy = λ_O
- 17) Wavelength of light in laboratory = λ_L
- 18) Current cosmic red shift = $z_{current} = z_c = z$
- 19) New cosmic red shift = $z_{new} = z_n = (z_c/1 + z_c) \leq 1$
- 20) Mean separation distance of any two neighboring galaxies = L_{sep}

Note: Planck scale symbols can be understood with a subscript 'pl' and current symbols can be understood with a subscript '0'.

1. INTRODUCTION

Modern cosmologists are seriously working on developing a peculiar model of relativistic cosmology that assumed to be started with a 'big bang' followed by a sudden and very short period exponential expansion called 'inflation' and a lately begun current acceleration. Big bang and inflation, both, seem to be the best widely accepted theoretical inferences of current large scale cosmic observations. But it is very clear to say that, both are having no proper theoretical

base and no way connected with the basic concepts of general theory of relativity. In addition to that, currently believed cosmic acceleration is based on another widely accepted invisible and unidentified cosmic 'new energy source' called 'dark energy'. Apart from these three complicated cosmic issues, galactic observations strongly supporting the existence of yet another mysterious and invisible cosmic 'mass source' called 'dark matter'. It is very unfortunate to say that, so far, no one did establish a clear physical connection between

cosmic dark energy and dark matter. Thus on a whole, currently believed standard cosmology is facing many challenges and many remarks. Since it is being developed and maintained by highly qualified scientists and high level mathematics, so called 'standard model' of cosmology is strongly believed and shown to be in line with large scale cosmic observations that are completely depending on an ad-hoc and misleading definition of 'cosmic red shift' associated with light coming from distant galaxies. Here we would like to emphasize the point that, by modifying the basic definition of cosmic red shift, considering 'speed of light' as an absolute cosmic expansion rate and adopting 'Planck mass' as the basic seed of the observed large scale universe, it is certainly possible to review and revise the basic picture of 'standard cosmology' and in near future, a perfect model of 'quantum cosmology' can be developed. To proceed further, in the following section, we are trying to highlight some important motivating points.

2. MOTIVATING POINTS

Even though standard model of cosmology is standing on 5 pillars namely, big bang, inflation, super luminal expansion, dark matter and dark energy, we would like to emphasize that,

1. We stress the point that - without a radial in-flow of matter in all directions towards one specific point, one cannot expect a big crunch and without a big crunch, one cannot expect a big bang. Really if there was a 'big bang' in the past, with reference to formation of big bang as supposed by cosmologists and with reference to the cosmic rate of expansion that might have taken place simultaneously in all directions at a 'naturally selected rate' about the point of big bang - 'point' of big bang can be considered as the characteristic reference point of cosmic expansion in all directions. Thinking in this way, to some extent, point of big bang can be considered as a possible centre of cosmic evolution. If so, thinking about a centre-less universe is illogical.
2. Even though highly intuitive and brilliant, concepts of big bang and inflation are no way connected with General relativity. It is a well known fact. In this context one can find interesting information in "<https://phys.org/news/2019-11-cosmologist-lonely-big-theory.html>".
3. Theoretically big bang, inflation, dark energy and super luminal expansion are no way connected with Planck scale which is having a major role in understanding quantum cosmology having information passing at speed of light.
4. Experimentally so far no one could understand big bang [1,2] and inflation [3,4], dark energy, dark matter and super luminal expansion with reference to any underground or ground or satellite based experiment.
5. Big bang, inflation and Super luminal expansion are no way giving a clue for unifying general theory of relativity and quantum mechanics.
6. Even though most of the cosmological observations are being studied and understood with photons that propagate with speed of light, it is very unfortunate to say that, most of the cosmologists are strongly believing in hypothetical 'super luminal expansion' of space. Recently detected gravitational waves that are supposed to originate from massive black holes are also confirmed to be moving at speed of light. If so, superluminal expansion can be considered as a pure human intellectual concept having no experimental support.
7. Big bang, inflation and dark energy are inference based intellectual concepts having no proper physical base and probably may misguide the future generation.
8. Compared to Big bang, inflation, dark energy and superluminal expansion, dark matter is having some sort of physical support in terms of an unknown, unidentified and unseen elementary particle having an heuristic gravitational attractive property. In addition to that, ultimately somehow, one should suppose the existence of some kind of matter for understanding the unexpected massive nature of trillions of observed galaxies.
9. 'Spin' is a basic property of quantum mechanics and 'rotation' is a very common experience.
10. It is better to understand and develop models of cosmology based on well supported physical concepts rather than extraordinary physical hypothesis that demand super-normal efforts.

3. OUR FIVE ASSUMPTIONS

We emphasize the point that, standard model of cosmology is basically a model of classical model of general relativity and based on the points proposed in 'Introduction', it seems inevitable to bring a change with reference to quantum model of cosmology. In this context, we propose five different assumptions in line with Planck mass as the baby universe. First three assumptions are helpful in understanding the basic cosmic structure and 4th and 5th assumptions are helpful in understanding galactic structures.

Right from the beginning of Planck scale, cosmic boundary is moving at speed of light and growing like a ball with the following workable assumptions [5-19].

Assumption-1: If $R_t \cong 2GM_t/c^2 \cong c/H_t$,
 $M_t \cong c^3/2GH_t$, and $tH_t \cong 1$.

Assumption-2: With reference to Planck mass, cosmic temperature follows a scaled form of Hawking's black hole temperature relation,
 $T_t \cong \frac{\hbar c^3}{8\pi Gk_B \sqrt{M_t M_{pl}}}$ where $M_{pl} \cong$ Planck mass.

Assumption-3: Ratio of Hubble parameter and angular velocity can be expressed as,
 $\frac{H_t}{\omega_t} \cong \Upsilon_t \cong 1 + \ln\left(\frac{H_{pl}}{H_t}\right)$ where H_{pl} is the Planck scale Hubble parameter.

Assumption-4: Galactic dark matter (M_{Gd})_t and visible matter (M_{Gv})_t are interrelated in such a way that,
 $M_{Gt} \cong (M_{Gd})_t + (M_{Gv})_t$ and

$$\frac{(M_{Gd})_t}{(M_{Gv})_t} \cong \sqrt{\frac{(M_{Gv})_t}{M_{Xt}}} \quad \text{where} \quad M_{Xt} \cong \frac{(M_t^3 M_{pl})^{\frac{1}{4}}}{8\pi} \cong$$

Time dependent dark-visible reference mass unit.

Assumption-5: Galactic flat rotation speed can be expressed as,
 $\frac{V_{Gt}}{c} \cong \left(\frac{1}{2\Upsilon_t}\right)^{\frac{1}{4}} \left(\frac{M_{Gt}}{M_t}\right)^{\frac{1}{4}}$ where
 $M_t =$ Cosmic total mass.

4. ESTIMATION OF THE MAGNITUDES OF PLANCK SCALE PARAMETERS

For the Planck scale, if, total mass = $M_{pl} \cong \sqrt{\frac{\hbar c}{G}}$, based on the assumptions, Planck scale physical parameters like, mass, radius, angular velocity, Hubble parameter and temperature can be estimated with the following relations.

$$R_{pl} \cong \frac{2GM_{pl}}{c^2} \cong 2\sqrt{\frac{G\hbar}{c^3}} \cong 3.2325 \times 10^{-35} \text{ m} \quad (1)$$

$$H_{pl} \cong \frac{c^3}{2GM_{pl}} \cong \frac{1}{2}\sqrt{\frac{c^5}{G\hbar}} \cong 9.27445 \times 10^{42} \text{ sec}^{-1} \quad (2)$$

$$t_{pl} \cong \frac{1}{H_{pl}} \cong 2\sqrt{\frac{G\hbar}{c^5}} \cong 1.07823 \times 10^{-43} \text{ sec} \quad (3)$$

$$\Upsilon_{pl} \cong \frac{H_{pl}}{\omega_{pl}} \cong 1 \quad (4)$$

$$T_{pl} \cong \frac{\hbar c^3}{8\pi Gk_B M_{pl}} \cong 5.6373 \times 10^{30} \text{ K} \quad (5)$$

$$\omega_{pl} \cong H_{pl} \cong 9.27445 \times 10^{42} \text{ rad.sec}^{-1} \quad (6)$$

5. ESTIMATION OF THE MAGNITUDES OF CURRENT COSMIC PARAMETERS

Considering, current cosmic microwave background radiation (CMBR) temperature [20] as, $T_0 \cong 2.725 \text{ K}$, current cosmic physical parameters like, mass, radius, angular velocity, Hubble parameter and galactic dark-visible reference mass unit can be estimated with the following relations.

$$M_0 \cong \frac{1}{M_{pl}} \left(\frac{\hbar c^3}{8\pi Gk_B T_0} \right)^2 \cong 9.31453 \times 10^{52} \text{ kg} \quad (7)$$

$$H_0 \cong \frac{c^3}{2GM_0} \cong 2.1671 \times 10^{-18} \text{ sec}^{-1} \cong 66.87 \text{ km.sec}^{-1} \text{ Mpc}^{-1} \quad (8)$$

$$\Upsilon_0 \cong 1 + \ln\left(\frac{H_{pl}}{H_0}\right) \cong 140.61 \quad (9)$$

$$\omega_0 \cong \frac{H_0}{\Upsilon_0} \cong \frac{2.1671 \times 10^{-18} \text{ sec}^{-1}}{140.61} \cong 1.5412 \times 10^{-20} \text{ rad.sec}^{-1} \quad (10)$$

$$R_0 \cong \frac{c}{H_0} \cong 1.3834 \times 10^{26} \text{ m} \quad (11)$$

$$t_0 \cong \frac{1}{H_0} \cong \frac{2GM_0}{c^3} \cong 14.62 \text{ Gy} \quad (12)$$

$$M_{X0} \cong \frac{(M_0^3 M_{pl})^{\frac{1}{4}}}{8\pi} \cong 3.623 \times 10^{38} \text{ kg} \quad (13)$$

6. THREE CHARACTERISTIC GALACTIC APPLICATIONS

Based on the assumptions, 3 characteristic properties of any galaxy can be expressed in the following way [16,17,18]. At present,

- 1) Radius of any galaxy can be expressed as [21,22]

$$R_G \cong \sqrt{\frac{GM_G}{c\omega_0}} \quad (14)$$

- 2) Angular velocity of any galaxy can be expressed as,

$$\omega_G \cong \frac{V_G^3}{GM_G} \quad (15)$$

- 3) Mean separation distance of any two neighboring galaxies can be expressed as,

$$L_{sep} \cong \left(\frac{\sqrt{V_{G1}V_{G2}}}{c} \right) \left(\frac{c}{H_0} \right) \cong \frac{\sqrt{V_{G1}V_{G2}}}{H_0} \quad (16)$$

where (V_{G1}, V_{G2}) represent the flat rotation speeds of galaxies $(G1, G2)$.

It may be noted that, according to Baryonic acoustic oscillations, current galactic separation distance is around 490 million light years [23,24]. In this context, relation (16) can be recommended for further research.

7. GALACTIC DARK MASS, VISIBLE MASS, TOTAL MASS, RADIUS AND ANGULAR VELOCITY

Based on the assumptions and observed galactic flat rotations speeds taken from Spitzer

Photometry and Accurate Rotation curves (SPARC) [25], in the following Table-1 we present the data for estimated galactic radii and angular velocities. Here it may be noted that, at present, for any galaxy, (ignoring the '0' subscript),

$$\begin{aligned} M_G &\cong M_{Gd} + M_{Gv} \\ &\cong M_{Gv} \sqrt{\frac{M_{Gv}}{M_{X0}}} + M_{Gv} \cong M_{Gv} \left(1 + \sqrt{\frac{M_{Gv}}{M_{X0}}} \right) \end{aligned} \quad (17)$$

$$\frac{V_G}{c} \cong \left(\frac{1}{2\Upsilon_0} \right)^{\frac{1}{4}} \left(\frac{M_G}{M_0} \right)^{\frac{1}{4}} \cong 0.2442 \left(\frac{M_G}{M_0} \right)^{\frac{1}{4}} \quad (18)$$

Replacing Modified Newtonian Dynamics (MOND) acceleration parameter with cosmic angular acceleration and replacing galactic visible mass with dark mass, relation (18) can be obtained. Considering a simple C++ program, relation (17) can be solved numerically. See Table 1. In most of the cases, estimated total mass of galaxy seems to be in line with galactic halo estimations made with Navarro–Frenk–White (NFW) model having marginal errors (http://astroweb.cwru.edu/SPARC/WP50_M200.mrt). It can be confirmed with columns (5), (6) of Table 1 and Fig. 1. Red curve is our estimation and green curve is for NFW model [26,27,28]. As total mass of galaxy is assumed to be proportional to 4th power of rotation speed, a small change in rotation speed will have large effect in galactic total mass. We have prepared Fig. 1 with respect to increasing galactic rotation speeds for a clear understanding. From the figure it is very clear that, for low rotation speeds, NFW estimated halo mass is on higher side and our estimated total mass is on lower side. Based on relation (14), if one is willing to consider the idea that, galactic radius is proportional to square root of its mass, it seems logical to say that, lower massive galaxies will have small radii and low rotation speeds. Here we would like appeal that, galaxies whose visible mass approaches our reference mass unit of $M_{X0} \cong 3.6 \times 10^{38} \text{ kg}$, seems to possess very little dark matter. It can be confirmed with very recent observations pertaining to NGC1052-DF2 and NGC1052-DF4 galaxies.

Recommended visible mass of NGC1052-DF2 [29,30,31] is $\approx 1 \times 10^8 M_{\text{Sun}}$ and its estimated dark mass is

$$\approx \left((2 \times 10^{38})^{\frac{3}{2}} / \sqrt{3.6 \times 10^{38}} \right) \cong 1.49 \times 10^{38} \text{ kg. Sum of}$$

dark mass and visible mass is $\approx 0.35 \times 10^9 M_{\text{Sun}}$. Corresponding flat rotation speed is 18.5 km/sec. It needs further study with respect to NGC1052-DF2 refined data. By means of tidal mass loss [32], if dark matter shifts from satellite galaxy to its mother galaxy, based on our proposed concepts, mother galaxy's flat rotation speed must increase significantly due to increase in total matter. It is for future observational testing.

8. UNDERSTANDING COSMIC RED SHIFT WITH SPEED OF LIGHT

It may be noted that, increased red shifts and increased distances forced Edwin Hubble to propose the Hubble's law [33]. With reference to laboratory, appropriate definition of red shift (z) seems to be [34].

$$z_{new} \cong z_n \cong \frac{\lambda_o - \lambda_L}{\lambda_o} \cong 1 - \left(\frac{\lambda_L}{\lambda_o} \right) \leq 1. \quad (19)$$

$$\text{But not } z_{current} \cong z_c \cong \frac{\lambda_o - \lambda_L}{\lambda_L} \cong \left(\frac{\lambda_o}{\lambda_L} \right) - 1 \quad (20)$$

Here, as usual, λ_o is the wave length of light received from observed galaxy and λ_L is the wave length of light in laboratory. Even though both relations are ad-hoc definitions, compared to relation (20), relation (19) seems to be appropriate with respect to 'light speed expansion'. Very interesting thing is that, when red shift is very small (up to $z \approx 0.01$), both relations almost all will give the same result. Important point to be noticed is that, by Hubble's time the maximum red shift noticed was 0.003.




Fig. 1. Comparison of estimated total mass of galaxy with NFW model

Table 1. Estimated galactic dark mass, visible mass, total mass, radius and angular velocity

Galaxy	Rotation speed (km/sec)	Error in rotation speed (km/sec)	Estimated total mass $1.989e40$ (kg)	Log10 (Total mass of galaxy/ Mass of Sun)	Galactic halo mass (NFW model)	Estimated visible mass $1.989e40$ (kg)	Estimated dark mass $1.989e40$ (kg)	Dark mass%	Estimated radius of galaxy (kpc)	Estimated angular velocity of galaxy (rad/sec)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
D631-7	57	9.4	1.72	10.24	10.7	0.33	1.39	80.94	22.79	8.11E-17
DDO064	46.1	3.9	0.74	9.87	10.53	0.18	0.56	75.78	14.91	1.00E-16
DDO154	47	1	0.80	9.90	10.58	0.19	0.61	76.30	15.49	9.83E-17
DDO161	66.3	1.9	3.15	10.50	10.87	0.50	2.65	84.02	30.83	6.97E-17
DDO168	53.4	1.9	1.33	10.12	10.67	0.27	1.05	79.46	20.00	8.65E-17
DDO170	60	1.6	2.11	10.32	10.65	0.38	1.73	82.04	25.25	7.70E-17
ESO079-G014	175	3.5	152.90	12.18	12.39	7.28	145.62	95.24	214.80	2.64E-17
ESO116-G012	109.1	3.1	23.10	11.36	11.49	2.01	21.09	91.30	83.49	4.24E-17
ESO563-G021	314.6	11.7	1596.97	13.20	13.45	35.41	1561.55	97.78	694.19	1.47E-17
F561-1	50	2.9	1.02	10.01	11.13	0.23	0.79	77.88	17.53	9.24E-17
F563-V2	116.6	9.4	30.13	11.48	11.12	2.41	27.73	92.00	95.36	3.96E-17
F568-V1	112.3	15.8	25.93	11.41	11.16	2.17	23.76	91.61	88.45	4.11E-17
F571-8	139.7	4.3	62.09	11.79	12.05	3.95	58.15	93.64	136.88	3.31E-17
F571-V1	83.6	3.5	7.96	10.90	10.98	0.96	7.00	87.91	49.02	5.53E-17
F574-1	97.8	4.1	14.91	11.17	11.27	1.49	13.43	90.03	67.09	4.72E-17
F579-V1	112.1	13.4	25.74	11.41	11.33	2.16	23.58	91.60	88.14	4.12E-17
F583-1	85.8	3.6	8.84	10.95	11.02	1.03	7.80	88.29	51.63	5.39E-17
IC2574	66.4	2	3.17	10.50	11.29	0.51	2.66	84.05	30.92	6.96E-17
IC4202	242.6	11	564.71	12.75	11.94	17.60	547.11	96.88	412.80	1.90E-17
KK98-251	33.7	1.6	0.21	9.32	10.34	0.07	0.14	66.38	7.97	1.37E-16
NGC0024	106.3	7.9	20.82	11.32	11.3	1.87	18.95	91.02	79.26	4.35E-17
NGC0055	85.6	5	8.75	10.94	11.27	1.03	7.73	88.25	51.39	5.40E-17
NGC0100	88.1	6.4	9.82	10.99	11.19	1.11	8.71	88.66	54.44	5.24E-17
NGC0247	104.9	8	19.74	11.30	11.37	1.80	17.94	90.87	77.18	4.40E-17
NGC0289	163	8	115.08	12.06	11.82	6.01	109.08	94.78	186.35	2.83E-17

Galaxy	Rotation speed (km/sec)	Error in rotation speed (km/sec)	Estimated total mass 1.989e40 (kg)	Log10 (Total mass of galaxy/ Mass of Sun)	Galactic halo mass (NFW model)	Estimated visible mass 1.989e40 (kg)	Estimated dark mass 1.989e40 (kg)	Dark mass%	Estimated radius of galaxy (kpc)	Estimated angular velocity of galaxy (rad/sec)
NGC0300	93.3	7	12.35	11.09	11.11	1.31	11.05	89.44	61.06	4.95E-17
NGC0801	220	16.2	381.90	12.58	12.32	13.52	368.39	96.46	339.47	2.10E-17
NGC0891	216.1	5.7	355.53	12.55	11.93	12.88	342.65	96.38	327.54	2.14E-17
NGC1003	109.8	4.2	23.70	11.37	11.59	2.04	21.65	91.37	84.56	4.21E-17
NGC1090	164.4	3.7	119.09	12.08	11.71	6.15	112.94	94.84	189.57	2.81E-17
NGC1705	71.9	4.3	4.36	10.64	10.82	0.63	3.73	85.49	36.26	6.43E-17
NGC2366	50.2	3.2	1.04	10.02	10.61	0.23	0.81	77.98	17.68	9.20E-17
NGC2403	131.2	4.9	48.31	11.68	11.4	3.33	44.98	93.11	120.73	3.52E-17
NGC2683	154	8.1	91.69	11.96	11.81	5.15	86.55	94.39	166.34	3.00E-17
NGC2841	284.8	8.6	1072.56	13.03	12.57	27.10	1045.46	97.47	568.91	1.62E-17
NGC2903	184.6	5.6	189.32	12.28	11.64	8.42	180.91	95.55	239.01	2.50E-17
NGC2915	83.5	6.3	7.93	10.90	10.82	0.96	6.97	87.89	48.90	5.53E-17
NGC2976	85.4	3.3	8.67	10.94	11.13	1.02	7.65	88.22	51.15	5.41E-17
NGC2998	209.9	8.1	316.45	12.50	12	11.91	304.55	96.24	309.02	2.20E-17
NGC3109	66.2	2.6	3.13	10.50	11.06	0.50	2.63	83.99	30.74	6.98E-17
NGC3198	150.1	3.9	82.75	11.92	11.75	4.80	77.95	94.20	158.02	3.08E-17
NGC3521	213.7	15.9	340.00	12.53	12.41	12.50	327.51	96.32	320.31	2.16E-17
NGC3726	168	6.2	129.87	12.11	11.93	6.52	123.35	94.98	197.96	2.75E-17
NGC3741	50.1	2.1	1.03	10.01	10.39	0.23	0.80	77.93	17.61	9.22E-17
NGC3769	118.6	8.4	32.26	11.51	11.42	2.53	29.73	92.17	98.66	3.90E-17
NGC3877	168.4	5.1	131.11	12.12	11.68	6.56	124.55	95.00	198.90	2.74E-17
NGC3893	174	8.9	149.44	12.17	11.93	7.17	142.27	95.20	212.35	2.66E-17
NGC3917	135.9	4.1	55.61	11.75	11.75	3.66	51.95	93.41	129.54	3.40E-17
NGC3949	163	7.1	115.08	12.06	11.71	6.01	109.08	94.78	186.35	2.83E-17
NGC3953	220.8	6.1	387.49	12.59	12.23	13.65	373.84	96.48	341.95	2.09E-17
NGC3972	132.7	2.9	50.55	11.70	11.57	3.43	47.12	93.21	123.51	3.48E-17
NGC3992	241	5.2	549.96	12.74	12.12	17.29	532.67	96.86	407.38	1.92E-17
NGC4010	125.8	4.7	40.83	11.61	11.6	2.97	37.86	92.73	111.00	3.67E-17

Galaxy	Rotation speed (km/sec)	Error in rotation speed (km/sec)	Estimated total mass $1.989e40$ (kg)	Log10 (Total mass of galaxy/ Mass of Sun)	Galactic halo mass (NFW model)	Estimated visible mass $1.989e40$ (kg)	Estimated dark mass $1.989e40$ (kg)	Dark mass%	Estimated radius of galaxy (kpc)	Estimated angular velocity of galaxy (rad/sec)
NGC4013	172.9	7.1	145.69	12.16	12.27	7.05	138.65	95.16	209.68	2.67E-17
NGC4051	157	5.5	99.05	12.00	11.92	5.43	93.63	94.52	172.89	2.94E-17
NGC4085	131.5	4.8	48.75	11.69	11.51	3.35	45.40	93.13	121.29	3.51E-17
NGC4088	171.7	6.9	141.69	12.15	11.94	6.92	134.78	95.12	206.78	2.69E-17
NGC4100	158.2	5	102.11	12.01	11.67	5.54	96.58	94.58	175.54	2.92E-17
NGC4138	147.3	5.9	76.75	11.89	11.83	4.56	72.19	94.06	152.18	3.14E-17
NGC4157	184.7	7.2	189.73	12.28	12.12	8.43	181.30	95.56	239.27	2.50E-17
NGC4183	110.6	5.4	24.39	11.39	11.22	2.09	22.31	91.45	85.80	4.18E-17
NGC4214	80.1	5.8	6.71	10.83	10.9	0.85	5.86	87.26	45.00	5.77E-17
NGC4217	181.3	7.2	176.14	12.25	12.15	8.01	168.13	95.45	230.55	2.55E-17
NGC4559	121.2	5.1	35.18	11.55	11.43	2.68	32.50	92.38	103.03	3.81E-17
NGC5005	262.2	20.7	770.53	12.89	12.86	21.70	748.84	97.18	482.20	1.76E-17
NGC5033	194.2	3.6	231.88	12.37	11.88	9.65	222.23	95.84	264.52	2.38E-17
NGC5055	179	4.9	167.37	12.22	11.72	7.74	159.63	95.37	224.73	2.58E-17
NGC5371	209.5	3.9	314.05	12.50	11.75	11.85	302.21	96.23	307.84	2.21E-17
NGC5585	90.3	2.4	10.84	11.04	11.26	1.19	9.65	89.00	57.19	5.12E-17
NGC5907	215	2.9	348.35	12.54	11.88	12.71	335.65	96.35	324.22	2.15E-17
NGC5985	293.6	8.6	1211.39	13.08	12.2	29.41	1181.99	97.57	604.61	1.57E-17
NGC6015	154.1	7	91.93	11.96	11.46	5.16	86.78	94.39	166.56	3.00E-17
NGC6195	251.7	9.3	654.33	12.82	12.73	19.44	634.89	97.03	444.35	1.84E-17
NGC6503	116.3	2.4	29.82	11.47	11.24	2.39	27.44	91.98	94.87	3.97E-17
NGC6674	241.3	4.9	552.70	12.74	12.8	17.35	535.36	96.86	408.39	1.91E-17
NGC6946	158.9	10.9	103.93	12.02	11.8	5.61	98.33	94.61	177.10	2.91E-17
NGC7331	239	5.4	531.93	12.73	12.42	16.91	515.03	96.82	400.64	1.93E-17
NGC7814	218.9	7	374.32	12.57	12.05	13.34	360.99	96.44	336.09	2.11E-17
PGC51017	18.6	1.3	0.02	8.29	10.25	0.01	0.01	43.91	2.43	2.48E-16
UGC00128	129.3	2.8	45.57	11.66	11.48	3.20	42.37	92.98	117.26	3.57E-17
UGC00731	73.3	2.3	4.71	10.67	10.72	0.67	4.04	85.82	37.69	6.30E-17

Galaxy	Rotation speed (km/sec)	Error in rotation speed (km/sec)	Estimated total mass 1.989e40 (kg)	Log10 (Total mass of galaxy/ Mass of Sun)	Galactic halo mass (NFW model)	Estimated visible mass 1.989e40 (kg)	Estimated dark mass 1.989e40 (kg)	Dark mass%	Estimated radius of galaxy (kpc)	Estimated angular velocity of galaxy (rad/sec)
UGC01230	103.7	6.1	18.85	11.28	11.28	1.75	17.11	90.73	75.43	4.46E-17
UGC01281	55.2	3.5	1.51	10.18	10.74	0.30	1.22	80.22	21.37	8.37E-17
UGC02259	86.2	2.9	9.00	10.95	10.94	1.05	7.95	88.35	52.12	5.36E-17
UGC02487	332	3.5	1980.67	13.30	12.26	40.92	1939.77	97.93	773.10	1.39E-17
UGC02885	289.5	12	1145.13	13.06	12.67	28.32	1116.82	97.53	587.84	1.60E-17
UGC02916	182.7	6.9	181.64	12.26	11.95	8.18	173.47	95.49	234.12	2.53E-17
UGC02953	264.9	6	802.77	12.90	12.23	22.30	780.47	97.22	492.18	1.74E-17
UGC03205	219.6	8.6	379.13	12.58	12.28	13.46	365.69	96.45	338.24	2.10E-17
UGC03546	196.9	7.4	245.04	12.39	11.89	10.02	235.03	95.91	271.93	2.35E-17
UGC03580	126.2	3.2	41.35	11.62	11.5	2.99	38.36	92.76	111.71	3.66E-17
UGC04278	91.4	4.8	11.38	11.06	11.16	1.23	10.15	89.16	58.59	5.06E-17
UGC04305	34.5	2.7	0.23	9.36	10.79	0.08	0.16	67.14	8.35	1.34E-16
UGC04325	90.9	2.7	11.13	11.05	11.03	1.21	9.92	89.09	57.95	5.08E-17
UGC04499	72.8	2.4	4.58	10.66	10.1	0.65	3.93	85.71	37.17	6.35E-17
UGC05005	98.9	7.2	15.60	11.19	11.1	1.53	14.07	90.17	68.60	4.67E-17
UGC05253	213.71	7	340.06	12.53	12.08	12.50	327.57	96.32	320.34	2.16E-17
UGC05716	73.1	1.2	4.66	10.67	10.77	0.66	4.00	85.78	37.48	6.32E-17
UGC05721	79.7	6.6	6.58	10.82	10.79	0.84	5.74	87.19	44.55	5.80E-17
UGC05986	113	4.1	26.58	11.42	11.71	2.21	24.37	91.68	89.56	4.09E-17
UGC06399	85	3.8	8.51	10.93	11.1	1.01	7.50	88.15	50.68	5.44E-17
UGC06446	82.2	4.3	7.44	10.87	10.89	0.92	6.53	87.66	47.39	5.62E-17
UGC06614	199.8	16	259.80	12.41	12.64	10.42	249.39	95.99	280.00	2.31E-17
UGC06628	41.8	6.4	0.50	9.70	11.11	0.13	0.36	73.08	12.26	1.11E-16
UGC06667	83.8	3.1	8.04	10.91	11.11	0.97	7.07	87.94	49.25	5.51E-17
UGC06786	219.4	7.8	377.75	12.58	12.08	13.42	364.33	96.45	337.62	2.11E-17
UGC06787	248.1	4.8	617.69	12.79	13.38	18.70	599.00	96.97	431.73	1.86E-17
UGC06818	71.2	4	4.19	10.62	10.9	0.62	3.57	85.32	35.56	6.49E-17
UGC06917	108.7	3.5	22.76	11.36	11.32	1.99	20.77	91.26	82.87	4.25E-17

Galaxy	Rotation speed (km/sec)	Error in rotation speed (km/sec)	Estimated total mass 1.989e40 (kg)	Log10 (Total mass of galaxy/ Mass of Sun)	Galactic halo mass (NFW model)	Estimated visible mass 1.989e40 (kg)	Estimated dark mass 1.989e40 (kg)	Dark mass%	Estimated radius of galaxy (kpc)	Estimated angular velocity of galaxy (rad/sec)
UGC06923	79.6	2.5	6.55	10.82	11.07	0.84	5.71	87.16	44.44	5.80E-17
UGC06930	107.2	5.1	21.53	11.33	11.3	1.91	19.62	91.11	80.60	4.31E-17
UGC06973	174.2	6.2	150.13	12.18	11.87	7.19	142.94	95.21	212.84	2.65E-17
UGC06983	109	5.8	23.01	11.36	11.23	2.00	21.01	91.29	83.33	4.24E-17
UGC07125	65.2	2.1	2.95	10.47	10.72	0.48	2.47	83.70	29.82	7.09E-17
UGC07151	73.5	2.8	4.76	10.68	11.07	0.67	4.09	85.87	37.89	6.29E-17
UGC07261	74.7	3.4	5.08	10.71	11	0.70	4.37	86.14	39.14	6.19E-17
UGC07399	103	3.3	18.35	11.26	11.12	1.71	16.64	90.66	74.41	4.49E-17
UGC07524	79.5	3.6	6.51	10.81	11.07	0.84	5.68	87.15	44.33	5.81E-17
UGC07603	61.6	2.8	2.35	10.37	10.77	0.41	1.94	82.58	26.61	7.50E-17
UGC07690	57.4	3.2	1.77	10.25	10.89	0.33	1.44	81.09	23.11	8.05E-17
UGC08286	82.4	2.3	7.52	10.88	10.87	0.93	6.59	87.69	47.62	5.61E-17
UGC08490	78.6	3.8	6.22	10.79	10.74	0.81	5.41	86.97	43.33	5.88E-17
UGC08550	56.9	1.9	1.71	10.23	10.65	0.33	1.38	80.90	22.71	8.12E-17
UGC08699	182.4	6.9	180.45	12.26	11.99	8.15	172.31	95.49	233.35	2.53E-17
UGC09037	152.3	9.6	87.71	11.94	12.09	5.00	82.72	94.31	162.69	3.03E-17
UGC09133	226.8	4.2	431.35	12.63	12.15	14.68	416.68	96.60	360.78	2.04E-17
UGC09992	33.6	3.3	0.21	9.32	10.64	0.07	0.14	66.25	7.92	1.38E-16
UGC10310	71.4	3.9	4.24	10.63	10.96	0.62	3.62	85.37	35.76	6.47E-17
UGC11455	269.4	7.4	858.72	12.93	13.03	23.34	835.39	97.28	509.04	1.72E-17
UGC11914	288.1	10.5	1123.14	13.05	13.59	27.96	1095.19	97.51	582.17	1.60E-17
UGC12506	234	16.8	488.79	12.69	12.08	15.97	472.83	96.73	384.05	1.97E-17
UGC12632	71.7	2.8	4.31	10.63	10.83	0.63	3.68	85.44	36.06	6.44E-17
UGCA444	37	4.8	0.31	9.49	10.07	0.09	0.21	69.39	9.60	1.25E-16

With reference to relation (20), relation (19) can be expressed as,

$$z_n \cong \frac{z_c}{1+z_c} \leq 1 \quad (21)$$

Based on this new definition, farthest galaxies distance can be estimated very easily. For example, see the following Table 2. We sincerely appeal that, on cosmological scales, 2.5% is not at all a 'serious' error. We would like to emphasize the point that, conceptually, we are no way deviating from the basic idea of expanding universe and receding galaxies. Only thing is that, we are confining to 'light speed expansion' and 'light speed receding'. With further study, there is a scope for understanding the universe in a unified approach. Since most of the cosmological observations are being studied with photons that move at speed of light, rather than 'working on controversial cosmic 'acceleration' and 'flatness' phenomena [35,15], it is better to work on understanding the root causes of 'speed of light'.

Richard Powell has written an online C program (<http://www.atlasoftheuniverse.com/cosmodis.c>) (version 1.1) for estimating the light travel distance. For a summary of the used formulae, one can refer David Hogg [36]. Using that program and considering a red shift of $z_c = (0.1$ to 200), we have prepared Figure-2. Green curve indicates the light travel distance in Lambda cosmology prepared with Omega matter = 0.32, Omega lambda = 0.68, Omega radiation = 0.0 and $H_0 = 66.87$ km/sec/Mpc. Red curve indicates our estimated light travel distance $[z_c/(1+z_c)](c/H_0) \cong [z_n](c/H_0)$ where $H_0 = 66.87$ km/sec/Mpc. As traditional red shift is increasing, error in estimated light travel distance is increasing to +8.59% at $z_c \cong 1.20$ and from there onwards, error is reaching to 0% at $z_c \cong 11.5$ to 11.55. Proceeding further, error is reaching to -5.14% at $z_c \cong 200.0$. Here, 'positive error' means, traditional light travel distance is higher than our estimate and 'negative error' means, traditional light travel distance is lower than our estimate. This can be also be confirmed with other online cosmic redshift-distance calculators written by Aaron Robotham and Joseph Dunne (<https://cosmocalc.icrar.org/>).

Based on the two curves presented in Figure-2, it is certainly possible to say that, if 'light travel distance' is a characteristic index in Lambda

cosmology, the same index, can also be understood with our re-defined red shift relation (19) and 'light speed expansion'. 'An interesting feature' is that the ratio of 'comoving distance' and 'light travel distance' is marginally and approximately equal to $\exp(z_n)$. We are working in this direction.

With reference to relation (21), relation (20) can be expressed as,

$$\text{Traditional redshift, } z_c \cong \frac{z_n}{1-z_n} \quad (22)$$

Relation (22) indicates that, for increasing light travel distances, z_c seems to have higher values compared to re-defined red shift z_n . See Figure 3. Red curve indicates our re-defined red shift and green curve indicates traditional red shift. To some extent, based on relations (21) and (22), it is possible to guess that, currently believed cosmic acceleration can be considered as an alternative interpretation associated with the alternative relation (20). It needs further study with respect to galactic red shifts and star rotation curves etc. Among relations (19) and (20), the correct definition can be decided with future observations.

9. UNDERSTANDING AND MODIFYING LAMBDA COSMOLOGY

With reference to traditional cosmology, understanding 'Lambda term' or 'cosmological constant' is a very difficult task. We would like to emphasize that, quantum cosmology point of either Lambda term or cosmological constant can be understood very easily. By considering Planck scale as the origin and growing Planck ball as a sequel of cosmic evolution, it is very simple to understand the cosmic physics. In this context, we appeal that,

- 1) Even though highly intuitive and impressive, there is no clarity and proper physical support for 'big bang' and 'inflation' concepts.
- 2) Lambda term can be expressed as a scalar quantity having the form,

$$\Lambda_t \cong \frac{3}{R_t^2} \cong \frac{3H_t^2}{c^2}.$$

- 3) Similarly, cosmological constant can be expressed as a scalar quantity having the

$$\text{form, } \frac{\Lambda_t c^4}{8\pi G}.$$

- 4) Based on the assumptions, total mass energy density and critical energy density, both are identical.
- 5) At any stage of cosmic evolution, $\frac{3c^2 H_t^2}{8\pi G} - \frac{\Lambda_t c^4}{8\pi G} \cong 0$. Clearly speaking, if one is willing to consider $\left(\frac{\Lambda_t c^4}{8\pi G}\right)$ as a characteristic expression for 'dark energy density', it can be inferred that, at any stage of cosmic evolution, difference of dark energy density and mass-energy density is always 'zero' and universe is always expanding at speed of light.
- 6) Frankly speaking, if universe is really expanding with 'speed of light', unphysical Lambda term and its inherited dark energy term, both can be relinquished forever. It needs further investigation.
- 7) In Lambda cosmology, matter creation is associated with big bang. Based on the assumptions, matter is being created continuously with an expression of the form, $M_t \cong \frac{c^3}{2GH_t} \cong \frac{c^3 t}{2G}$.
- 8) Cosmic Temperature-Time relation can be expressed as, $T_t \cong \frac{0.18511 \times 10^{10}}{\sqrt{t}}$ K. This relation is almost similar to the mainstream relation derived on big bang concepts. Only difference is that, for the same expected temperatures, our estimated cosmic physical processes are taking place early compared to big bang model. This idea helps in understanding the early formation of galaxies at the cosmic dawn.
- 9) Most important thing is that the characteristic cosmic expansion rate can be accurately estimated by knowing the CMBR temperature and there is no need to take the help of galactic red shifts and galactic distances.
- 10) Estimated galactic dark matter is more than 90 percent of the total mass of 'massive' galaxies. In case of least massive galaxies also, dark matter is roughly 50% of the total mass of galaxy.
- 11) Advanced galactic red shift data is raising doubts on well believed cosmic acceleration [37,38] and supporting constant rate of expansion [39]. Considering the Tolman test [40] for surface brightness and based on the analysis of the UV SB of luminous disk galaxies from HUDF and GALEX datasets, reaching from the local universe to $z \sim 5$, recently it has been shown that the surface brightness remains constant as expected in a static universe [41,42].
- 12) One thing is very clear that, red shift data analysis associated with cosmic rate of expansion is generating lot of confusion and controversy in assessing the correct rate of cosmic expansion. In this context, based on the data presented in Table 2, our proposed definition of red shift i.e. relations (20) and (22) can be recommended for further research.
- 13) Quantum cosmology point of view, Lambda cosmology needs a review at fundamental level. Our assumptions are based on 'time reversed' black holes and seem to be well connected with General theory of relativity as well as Quantum mechanics [5,6,13].
- 14) By counting the actual number of galaxies and considering the average mass of galaxy and with minor adjustments, our assumptions can be refined in workable way compared to Lambda model of cosmology.
- 15) In 2011, researchers found evidence that galaxies tend to rotate in a preferred direction. They uncovered an excess of left-handed, or counter-clockwise rotating, spirals in the part of the sky toward the north pole of the Milky Way. This study suggests that the shape of the big bang might be more complicated than the previously thought and that the early universe spun on an axis [43]. Recently, research presented at the 236th meeting of the American Astronomical Society suggests that the whole universe may be spinning. If further studies bear this out, the finding will challenge some of the fundamental assumptions of modern cosmology [44]. In this context, considering the field experts [45,46,47,48,49] advocated value of 'current cosmic angular velocity $\approx (10^{-13} \text{ rad}\cdot\text{year}^{-1})$ ', our assumptions can be recommended for observational search.
- 16) When it is assumed that, current universe is a rotating white hole, there exists no scope for causal disconnection of galaxies and no scope for 'horizon' problem. Similarly, when it is assumed that, universe is a rotating white hole having a positive curvature and expanding/growing with speed of light, then 'light speed expanding space' can be called as 'Flat space'.

Table 2. Estimation and fitting of the distances of farthest galaxies with re-defined red shift

Galaxy	Red shift	Standard Light travel distance (Gly)	Estimated Light travel distance (Gly)	%Error
GN-z11	11.09	13.39	13.41	-0.15
MACS1149-JD1	9.11	13.26	13.17	0.65
EGSY8p7	8.68	13.23	13.11	0.91
A2744 YD4	8.38	13.2	13.06	1.05
EGS-zs8-1	7.73	13.13	12.95	1.41
z7 GSD 3811	7.66	13.11	12.93	1.36
z8_GND_5296	7.51	13.1	12.9	1.51
SXDF-NB1006-2	7.215	13.17	12.84	2.5
GN-108036	7.213	13.07	12.84	2.5
BDF-3299	7.109	13.05	12.84	2.5
A1703 zD6	7.014	13.04	12.84	2.5
BDF-521	7.008	13.04	12.84	2.5
G2-1408	6.972	13.03	12.84	2.5
IOK-1	6.964	13.03	12.84	2.5




Fig. 2. Comparison of standard and estimated light travel distances




Fig. 3. Comparison of standard and re-defined red shifts

10. TO DERIVE COSMIC TEMPERATURE RELATION

At any stage of cosmic evolution and thermal expansion, with reference to Planck mass $M_{pl} \cong \sqrt{\hbar c/G}$ and by considering growing cosmic mass M_t as a point particle separated by an increasing hypothetical distance equal to $r_t \cong$ Wiens's thermal wavelength/ 2π , gravitational force of attraction can always be shown to be approximately equal to Einstein's force constant ($c^4/8\pi G$). Beauty of this procedure is that in a microscopic view, it integrates Newton's law of gravitation, Einstein's cosmic force constant, Wien's displacement law, Wave-Particle relation, Special theory of relativity and Hawking's black hole temperature formula.

Hypothesis-1: Considering universe as a point particle of mass M_t and gravitational force of attraction between the point universe and the Planck mass (the baby universe mass) is,

$$\frac{GM_t M_{pl}}{r_t^2} \cong \frac{c^4}{8\pi G} \quad (23)$$

where r_t is the hypothetical distance between (M_t, M_{pl}) .

Hypothesis-2: Hypothetical distance between point universe and Planck mass is always equal to,

$$r_t \cong \frac{\text{Wien's thermal wave length}}{2\pi} \cong \frac{1}{2\pi} \left(\frac{2.898 \times 10^{-3}}{T_t} \right) \quad (24)$$

Hypothesis-3: Mass of universe is always equal to,

$$M_t \cong \frac{c^3}{2GH_t} \quad (25)$$

Based on these three hypotheses,

$$T_t \cong \frac{1}{\sqrt{8\pi * 4.96511422^2} k_B G \sqrt{M_t M_{pl}}} \cong \frac{\hbar c^3}{24.891 k_B G \sqrt{M_t M_{pl}}} \quad (26)$$

Comparing this result with our assumption-3, with 0.96% error,

$$T_t \cong \frac{\hbar c^3}{24.891 k_B G \sqrt{M_t M_{pl}}} \cong \frac{\hbar c^3}{8\pi k_B G \sqrt{M_t M_{pl}}} \quad (27)$$

We have,

$$\left(\sqrt{8\pi * 4.965^2} \cong 24.8914 \right) \square \left(8\pi \cong 25.13274123 \right) \quad (28)$$

For the current case, if $T_0 \cong 2.725$ K,

$$M_0 \cong \left(\frac{\hbar c^3}{24.8914 k_B T_0 \sqrt{M_{pl}}} \right)^2 \cong \left(\frac{8.39726 \times 10^{26}}{2.725} \right)^2 \cong 9.496 \times 10^{52} \text{ kg} \quad (29)$$

$$H_0 \cong \frac{c^3}{2GM_0} \cong \left(\frac{2.01856 \times 10^{35}}{M_0} \right) \text{sec}^{-1} \quad (30)$$

$$\cong 2.1257 \times 10^{-18} \text{ sec}^{-1} \cong 65.59 \text{ km/sec/Mpc}$$

11. APPLICATIONS AND MORE FUTURE PROSPECTS

We would like to appeal that,

A) Considering the new definition of cosmic red shift,

- 1) Confusion associated with basic cosmic red shift can be eliminated.
- 2) Galactic flat rotation speeds and their distances can be reviewed in a better way.
- 3) Complicated mathematical formula associated with light travel distances can be simplified.

B) Considering Planck mass and light speed expansion

- 1) Standard cosmology and basic models of quantum cosmology can be studied in a unified approach.
- 2) Big bang, inflation and dark energy like intellectual concepts can be explored at utmost fundamental level.
- 3) Complicated mathematics of general theory of relativity can be simplified to a great extent.
- 4) Planck scale imprints associated with current cosmic observations can be explored and thereby a correct model of cosmology can be developed with high precision.

C) Considering the new formula associated with dark matter and visible matter

- 1) Inter relation between galactic dark matter and visible matter can be refined and understood in a better way.

- 2) Nature of dark matter can be explored at utmost fundamental level.
- 3) MOND scheme can be reviewed for a better understanding.
- 4) Atomic and nuclear scale dark matter secrets can be understood and thereby experiments can be conducted in a result oriented approach.

D) Considering the proposed cosmic angular velocity relation

- 1) Inter relation between actual cosmic expansion rate and actual cosmic angular velocity can be understood in a better way [50].
- 2) Ground reality of cosmic rotation can be verified with other cosmic observations.
- 3) Galactic radii and angular velocity relations can be refined with respect to actual galactic mass.

12. CONCLUSION

When modern cosmology was in its developing stage, black hole physics was in budding stage. Now, black hole physics is in its advanced stage and modern Lambda cosmology is facing many challenges from all corners. Another important point to be noted is that, when modern cosmology is in its developing stage, subject of dark matter is mostly in its controversial stage. Considering our five assumptions, to the possible extent, subject of cosmology, galactic dark matter, galactic flat rotations curves and galactic radii can be studied in a unified manner. Considering assumptions (1) and (2), tension in estimating the value of current Hubble constant can be eliminated in a unified approach with great confidence. Clearly speaking, first two assumptions play a vital role in understanding the basic structure of the expanding universe in a very simplified quantum gravitational approach independent of currently believed classical model of Lambda cosmology. Another noteworthy point is that in reality, it is not at all possible to reach any black hole, but the idea of – ‘observers residing in a time-reversed black hole (a white hole) universe that follows a scaled Hawking’s temperature formula - is quite exciting and paving a way for building a realistic model of quantum cosmology. Considering assumption (3), current cosmic angular velocity can be estimated and it can be understood and realized with the works of field experts like Godel, Birch, Godlowski, Longo, Chechin, Yuri Obukhov & team and Shamir. We are working on the non

equality of current Hubble parameter and current angular velocity and their backend physics. Considering assumption (4), it is possible to say that, role of dark matter in solar system and nuclear physics is negligible and it is observationally true. Considering assumption (5), galactic rotation speeds and total masses can be fitted well and there is a scope for applying it to other galaxies.

We are very confident to say that, along with our new red shift definition, our first three assumptions play a vital role in understanding the observed Universe ‘as a whole’. In addition to that, big bang and inflation like non-general relativistic concepts can be relinquished with further study. Proposed red shift definition and estimated method of galactic light travel distance is very simple and clearly indicates that, currently believed matter density fractions need not be required in developing a real model of cosmology. 4th and 5th assumptions help in quantifying the galactic dark matter.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

DATA AVAILABILITY STATEMENT

- 1) The data underlying this article pertaining to galaxies flat rotation speeds are available at <http://astroweb.cwru.edu/SPARC/>. It is being maintained by Federico Lelli, Stacy McGaugh and James Schombert.
- 2) The dataset underlying this article pertaining to farthest galaxies has been taken from https://en.wikipedia.org/wiki/List_of_the_most_distant_astronomical_objects.
- 3) Dataset C program underlying this article pertaining to light travel distances and red shift has been taken from <http://www.atlasoftheuniverse.com/cosmodis.c>. It was written by Richard Powell. To confirm our estimations, we have also considered cosmic redshift-distance

calculator written by Aaron Robotham and Joseph Dunne. Its URL is: <https://cosmocalc.icrar.org/>.

ACKNOWLEDGEMENTS

Author Seshavatharam is indebted to Professors Shri M. Nagaphani Sarma, Chairman, Shri K.V. Krishna Murthy, founder Chairman, Institute of Scientific Research in Vedas (I-SERVE), Hyderabad, India and Shri K.V.R.S. Murthy, former scientist IICT (CSIR), Govt. of India, Director, Research and Development, I-SERVE, for their valuable guidance and great support in developing this subject. Dr. Tatum thanks Dr. Rudolph Schild of the Harvard-Smithsonian Center for Astrophysics for his past support and encouragement.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Abhas Mitra. Why the big bang model cannot describe the observed universe having pressure and radiation. *Journal of Modern Physics*. 2011;2:1436.
2. Abhas Mitra. Why the Big Bang Model does not allow inflationary and cyclic cosmologies though mathematically one can obtain any model with favourable assumptions. *New Astronomy*. 2014;30:46
3. Roger Penrose. Difficulties with inflationary cosmology. *Annals of the New York Academy of Sciences*.1989;571:249.
4. Steinhardt PJ, Abraham Loeb. Inflationary schism. *Phys Lett B*. 2014;736:142
5. Roger Penrose. Gravitational collapse and space-time singularities. *Phys Rev Lett*. 1965;14:57.
6. Hawking S, Penrose R. The singularities of gravitational collapse and cosmology. *proc. roy. Soc Lond. A*. 1970;314:529-548
7. Hawking SW. Particle creation by black holes S. W. Hawking. *Communications in Mathematical Physics*. 1975;43(199):314-529.
8. Milgrom M. A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *ApJ. Part 1*. 1983;270:365.
9. Brownstein JR, Moffat JW. Galaxy rotation curves without non-baryonic dark matter. *ApJ*. 2006;636:721.
10. Melia F, Shevchuk ASH. The $R_h=ct$ universe. *MNRAS*. 2012;419:2579.
11. Tatum ET, Seshavatharam UVS, Lakshminarayana S. Flat space cosmology as a mathematical model of quantum gravity or quantum cosmology. *International Journal of Astronomy and Astrophysics*. 2015;5:133.
12. Tatum ET, Seshavatharam UVS, Lakshminarayana S. The basics of flat space cosmology. *International Journal of Astronomy and Astrophysics*. 2015;5:116.
13. Seshavatharam UVS, Tatum ET, Lakshminarayana S. On the role of gravitational self energy density in spherical flat space quantum cosmology. *Journal of Applied Physical Science International*. 2015;4(4):228.
14. Tatum ET, Seshavatharam UVS. How a Realistic Linear $R_h=ct$ Model of Cosmology Could Present the Illusion of Late Cosmic Acceleration. *Journal of Modern Physics*. 2018;9:1397.
15. Moncy John V. $R_h = ct$ and the eternal coasting cosmological model. *MNRAS*, 2019;484:L35.
16. Seshavatharam UVS, Lakshminarayana S. Light speed expansion and rotation of a primordial cosmic black hole universe having internal acceleration. *International Astronomy and Astrophysics Research Journal*. 2020;2(2):9.
17. Seshavatharam UVS, Lakshminarayana S. To correlate galactic dark and visible masses and to fit flat rotation speeds via monod approach and cosmic angular acceleration. *International Astronomy and Astrophysics Research Journal*. 2020; 2(3):28.
18. Seshavatharam UVS, Lakshminarayana S. Light speed expansion and rotation of a very dark machian universe having internal acceleration. *International Journal of Astronomy and Astrophysics*. 2020;10: 247.
19. Stacy McGaugh. Predictions and outcomes for the dynamics of rotating galaxies. *Galaxies*. 2020;8(2):35.
20. Planck Collaboration: Planck Results. *Cosmological Parameters*. 2015;13.

21. Laura Watkins L, et al. Evidence for an Intermediate-Mass Milky Way from Gaia DR2 Halo Globular Cluster Motions. *ApJ*, 2019;873:118.
22. Alis J, Deason et al. The edge of the galaxy. *MNRS*, 2020;496(3):3929.
23. Eisenstein DJ, et al. Detection of the baryon acoustic peak in the large-scale correlation function of SDSS luminous red galaxies. *Astrophys J*. 2005; 633:560.
24. César Hernández-Aguayo et al. Measuring the BAO peak position with different galaxy selections. *MNRS*. 2020;494(3):3120.
25. Federico Lelli, Stacy S, Mc Gaugh, James. Schombert M. SPARC: Mass models for 175 disk galaxies with spitzer photometry and accurate rotation curves. *ApJ*. 2016;152:6.
26. Navarro, Julio F, Frenk, Carlos S, White Simon DM. The structure of cold dark matter halos. *The Astrophysical Journal*. *ApJ*. 1996;462:563.
27. Pieter van Dokkum et al. A second galaxy missing dark matter in the NGC1052 group. *Ap Let*. 2019;874:L5.
28. Oman Kyle, et al. The unexpected diversity of dwarf galaxy rotation curves. *MNRS*, 2015;452(4):3650.
29. Shany Daniel, et al. Still missing dark matter: KCWI high-resolution stellar kinematics of NGC1052-DF2. *ApJ Let*. 2019;874:L12.
30. Pengfei Li, et al. A comprehensive catalog of dark matter halo models for SPARC galaxies. *ApJS*. 2020;247:31.
31. Lewis Geraint F et al. The globular cluster population of NGC 1052-DF2: Evidence for rotation. *MNRS Lett*. 2020;495(1):L1.
32. Daneng Yang, Hai-Bo Yu, Haipeng An. Self-interacting dark matter and the origin of ultradiffuse galaxies NGC1052-DF2 and -DF4. *Phys Rev Lett*. 2020;125:111105.
33. Hubble EP. A relation between distance and radial velocity among extra-galactic nebulae. *PNAS*. 1929;15:168.
34. Seshavatharam UVS, Lakshminarayana S. Basic interactions in black hole cosmology. *American Journal of Astronomy and Astrophysics*. 2013;2(1):6-17.
35. Abhas Mitra. Why the $R_h = ct$ cosmology is unphysical and in fact a vacuum in disguise like the milne cosmology. *MNRS*. 2014;442(1):382.
36. David W Hogg. Distance measures in cosmology. preprint (arXiv:9905116); 2000.
37. Riess AG, et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*. 1998;116:1009.
38. Perlmutter S, et al. Measurements of Ω and Λ from 42 high-redshift supernovae. *ApJ*. 1999;517:565.
39. Nielsen JT, Guffanti A, Sarkar S. Marginal evidence for cosmic acceleration from type ia supernovae. *Scientific Reports*, 2016;6: 35596
40. Tolman RC. On the estimation of distances in a curved universe with a non-static line element. *Proc. Natl. Acad Sci*. 1930; 16:511.
41. Eric Lerner J, Renato Falomo, Riccardo Scarpa. UV surface brightness of galaxies from the local universe to $z \sim 5$. *IJMPD*. 2014;23(6):1450058.
42. Eric J Lerner. Observations contradict galaxy size and surface brightness predictions that are based on the expanding universe hypothesis. *MNRS*. 2018;477(3):3185.
43. Michael Longo J. Detection of a dipole in the handedness of spiral galaxies with redshifts $z \sim 0.04$. *Phy. Let. B*. 2011;699(4):224.
44. Shamir L. Patterns of galaxy spin directions in SDSS and Pan-STARRS show parity violation and multipoles. *Astrophys Space Sci*. 2020;365:136.
45. Godel K. Rotating universes in general relativity theory. *Proceedings of the International Congress of Mathematicians in Cambridge*. 1950;1:175-181.
46. Birch P. Is the Universe rotating? *Nature*, 1982;298:451.
47. Godlowski W. Global and local effects of rotation: observational aspects. *IJMPD*. 2011;20:1643.
48. Chechin LM. Does the cosmological principle exist in the rotating universe? *Gravitation and Cosmology*. 2017;23:305.
49. Vladimir Korotky A, Eduard Masár Yuri Obukhov N. In the quest for cosmic rotation. *Universe*. 2020;6:14.

50. Sivaram Chandra, Kenath Arun. Primordial rotation of the universe, hydrodynamics, vortices and angular momenta of celestial objects. The Open Astronomy Journal. 2012;5:7.

© 2021 Seshavatharam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/66092>